

HUBBLE SPACE TELESCOPE WIDE FIELD CAMERA 3 OBSERVATIONS OF ESCAPING LYMAN CONTINUUM RADIATION FROM GALAXIES AND WEAK AGN AT REDSHIFTS $z \simeq 2.3\text{--}5$.

BRENT M. SMITH¹, ROGIER A. WINDHORST¹, ROLF A. JANSEN¹, SETH H. COHEN¹, LINHUA JIANG^{1,2}, MARK DIJKSTRA³,
ANTON M. KOEKEMOER⁴, RICHARD BIELBY⁵, AKIO K. INOUE⁶, JOHN W. MACKENTY⁴, ROBERT W. O'CONNELL⁷, JOSEPH I. SILK⁸

¹School of Earth & Space Exploration, Arizona State University, Tempe, AZ 85287-1404, USA; e-mail: Brent.Smith.1@asu.edu

²Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, P. R. China

³Institute of Theoretical Astrophysics, University of Oslo, Postboks 1029, 0858 Oslo, Norway

⁴Space Telescope Science Institute, Baltimore, MD 21218, USA

⁵University of Durham, South Road, Durham DH1 3LE, UK

⁶Osaka Sangyo University, Osaka, Japan

⁷Department of Astronomy, University of Virginia, Charlottesville, VA 22904-4325, USA

⁸The Johns Hopkins University, Baltimore, MD 21218, USA

Abstract

We present observations of escaping Lyman Continuum (LyC) radiation from 50 massive star-forming galaxies and 14 weak AGN with reliably measured spectroscopic redshifts at $z \simeq 2.3\text{--}5.8$. We analyzed Hubble Space Telescope (*HST*) Wide Field Camera 3 (WFC3) mosaics of the Early Release Science (ERS) field in three UVIS filters, and optical data from the Advanced Camera for Surveys (ACS) of the GOODS-South field to sample the rest-frame LyC over this redshift range. With our best current assessment of the WFC3 and ACS systematics, we find that the average LyC emission of galaxies at $\langle z \rangle = 2.38, 2.68, 3.47$, and 5.02 is generally measured at the $\gtrsim 3\sigma$ level in image stacks of 11–15 galaxies in the WFC3/UVIS F225W, F275W, F336W, and ACS/WFC F435W filters, respectively. Their measured average LyC flux correspond to $AB \simeq 29.5\text{--}30.7$ mag. The LyC flux of weak AGN is also detected to be typically ~ 1 mag brighter at $z \simeq 2.3\text{--}4.8$, but averaged over $\sim 4\times$ fewer galaxies in total. The stacked galaxy LyC light-profiles are flatter than their corresponding non-ionizing UV-continuum profiles out to radii of $r \lesssim 0''.7$, which may indicate a radial dependence of porosity in the ISM. The average LyC emission from AGN is more extended and sometimes more elongated compared to galaxies without AGN, which may be due to the apparent angle at which LyC escapes from AGN. With synthetic stellar SEDs fit to UV-continuum measurements longwards of Ly α and IGM transmission models, we find that the observed LyC flux most likely corresponds to absolute LyC escape fractions of $f_{\text{esc}}^{\text{abs}} \simeq 0.1\text{--}18\%$ at $z \simeq 2.3\text{--}3.9$, then increasing to a much wider range at $\langle z \rangle = 5.02$. All available data for galaxies, including published work, suggests a more sudden increase of f_{esc} with redshift at $z \simeq 3$. Our models imply that dust accumulating in (massive) galaxies over cosmic time may be a major factor in reducing f_{esc} more suddenly at $z \lesssim 3$. Our best fits of $f_{\text{esc}}(z)$ suggest that star-forming galaxies collectively contributed to maintaining cosmic reionization at redshifts $z \gtrsim 3\text{--}6$, while AGN likely dominated reionization at $z \lesssim 2.5$.

Keywords: Faint galaxies — UV imaging — Lyman-continuum — escape fractions.

1. INTRODUCTION

At the end of the cosmic dark ages, radiation emitted by the first objects that shined in the universe began to reionize the intergalactic medium (IGM). The far-ultraviolet (FUV) ionizing radiation, and specifically Lyman-continuum (“LyC”; $\lambda \leq 912\text{\AA}$), emitted by massive stars in the first star-forming galaxies (SFGs), or accretion disks around super-massive black holes in early Active Galactic Nuclei (AGN), may have initiated the epoch of cosmic reionization (e.g., [Madau et al. 2004](#)). Additional sources of LyC radiation and high energy particles within galaxies, such as high-mass X-ray binaries, galactic outflows/inflows and superwinds, accretion onto dark matter halos, massive pre-galactic Pop III stars, and young globular clusters may have contributed to the reionization of the IGM as well (e.g., [Ricotti 2002](#); [Sternberg et al. 2002](#);

[Mirabel et al. 2011](#); [Kulkarni et al. 2014](#)). This LyC radiation would have formed bubbles of ionized hydrogen around these UV-bright galaxies, which then expanded and merged until the IGM became completely ionized (e.g., [Gnedin 2000](#); [MiraldaEscudé et al. 2000](#); [Loeb & Barkana 2001](#); [Fan et al. 2002](#)). This phase transition of the neutral IGM began somewhere in the epoch $z \simeq 10\text{--}20$ ([Hinshaw et al. 2013](#); [Ade et al. 2014](#); [Planck Collaboration 2015](#)), and completed when the IGM was fully ionized by $z \simeq 6$ ([Mesinger & Haiman 2004](#); [Fan et al. 2006a, 2006b](#); [Schroeder et al. 2012](#); [McGreer et al. 2014](#); [Becker et al. 2015](#)), with observations of Ly α emitting galaxies favoring (volume-averaged) neutral fractions in excess of $\langle x_{\text{HI}} \rangle > 0.3$ at $z \sim 7$ (e.g., [Dijkstra et al. 2011](#); [Jensen et al. 2012](#); [Mesinger et al. 2014](#); [Choudhury et al. 2015](#)).

Because neutral hydrogen and dust are opaque to FUV radi-

ation, LyC photons can only escape from galaxies in regions where the surrounding H I column density, N_{H} , and dust extinction are low. Thus, in order for a fraction of the produced LyC photons to escape (f_{esc}), the interstellar medium (ISM) in the galaxy and its surrounding circumgalactic medium must be cleared. This can be accomplished by supernova winds (Fujita et al. 2003), which can also suppress the formation of low-mass stars and increase the formation of LyC producing high-mass stars, and can be further enhanced by AGN outflows (Silk & Norman 2009). High star-formation rates can also increase the porosity of the ISM (Clarke & Oey 2002). Semi-analytical models of Dove et al. (2000) show that LyC emitted by OB associations can become trapped in superbubbles until they expand outside of the disk. Once the surrounding medium is either cleared or fully ionized, it becomes transparent to LyC radiation, which can then escape through these regions of the galaxy, or be Thomson scattered by free electrons and/or dust. The escaping LyC can then be observed along some lines-of-sight, which can be distributed randomly in a galaxy, and is in some cases offset from the galactic center (e.g., Gnedin 2008; Iwata et al. 2009; Vanzella et al. 2010b, 2012).

Stacks of ground-based spectra have shown that AGN produce more LyC than star-forming Lyman Break Galaxies (LBG; Vanden Berk et al. 2001; Shapley et al. 2003), though LBGs selected via drop-out techniques may have fainter LyC emission due to their selection compared to other UV-bright SFGs (Vanzella et al. 2015). Rest-frame UV spectra of AGN taken with *HST* and the Far Ultraviolet Spectroscopic Explorer (*FUSE*; Moos et al. 2000) have shown significant detections of escaping LyC flux at $0.5 \lesssim z \lesssim 2.5$ (e.g., Telfer et al. 2002; Scott et al. 2004; Shull et al. 2012; Lusso et al. 2015), but only upper limits of $f_{\text{esc}} \lesssim 1\text{--}2\%$ from galaxies at the same redshifts (e.g., Bridge et al. 2010; Siana et al. 2010; Rutkowski et al. 2015; Sandberg et al. 2015). AGN contributed the majority of LyC photons to the ionizing background from their peak epoch at $z \simeq 2$ until today, and maintain the ionized state of the IGM (Cowie et al. 2009). However, because AGN are much more rare than galaxies, and their space density decreases at $z \gtrsim 2$ (Silverman et al. 2008; Ebrero et al. 2009; Aird et al. 2015), AGN likely did not reionize the IGM at $z \gtrsim 3$ (Madau 1999; Fan et al. 2001; Hunt et al. 2004), though they are believed to be the only sources responsible for He II reionization at $z \simeq 3$ (Haardt & Madau 2012; Worseck et al. 2014). Therefore, SFGs are regarded as the most likely candidates that started the reionization of the IGM at $z \gtrsim 6$ (but see, e.g., Madau & Haardt 2015).

Since higher IGM opacity at $z \gtrsim 6$ prevents a direct study of LyC emission from SFGs at this epoch, we must study lower redshift analogs in order to understand the sources of reionization of the IGM. Despite many attempts, rest-frame far-UV observations of SFGs at $0.5 \lesssim z \lesssim 2.5$ have so far not yielded significant detections of escaping LyC flux (e.g., Ferguson 2001; Giallongo et al. 2002; Fernández-Soto et al. 2003; Malkan et al. 2003; Inoue et al. 2005; Siana et al. 2007, 2010; Cowie et al. 2009; Bridge et al. 2010; Grazian et al. 2016; Rutkowski et al. 2015; Sandberg et al. 2015; Guaita et al. 2016). Ground-based spectra (Steidel et al. 2001; Shapley

et al. 2006; Cooke et al. 2014; de Barros et al. 2015) and optical narrow-band and broadband imaging of SFGs at $3 \lesssim z \lesssim 4$ (Iwata et al. 2009; Vanzella et al. 2010b, 2012; Boutsia et al. 2011; Nestor et al. 2011, 2013; Mostardi et al. 2013) have revealed evidence for escaping LyC photons along several sight-lines, with $f_{\text{esc}} \simeq 1\text{--}40\%$ despite higher IGM opacities at these higher redshifts (Haardt & Madau 1996, 2012). Furthermore, Vanzella et al. (2012) estimate f_{esc} for one LBG (GDS J033216.64–274253.3 at $z=3.795$) to be $>25\%$, although some ground-based measurements of escaping LyC may be contaminated with non-ionizing flux from blended lower redshift foreground interlopers due to the lower resolution of ground-based seeing (Vanzella et al. 2010a; Nestor et al. 2013; Mostardi et al. 2015; Siana et al. 2015). Spectroscopy of gamma-ray burst afterglows from $2 < z < 8$ have also been used to constrain f_{esc} to $<6\%$ at these redshifts (Chen et al. 2007; Fynbo et al. 2009; Wyithe et al. 2010).

Observations of some local starburst galaxies have shown significant, yet varying f_{esc} values (Leitherer et al. 1995; Hurwitz et al. 1997; Tumlinson et al. 1999; Deharveng et al. 2001; Heckman et al. 2001; Borthakur et al. 2014; Izotov et al. 2016), although Hanish et al. (2010) find that local starburst galaxies do not exhibit higher escape fractions compared to ordinary local SFGs. H II regions in nearby galaxies have been observed to release 40–75% of the LyC photons produced by massive stars into the local IGM (Ferguson et al. 1996; Leitherer et al. 1996; Oey & Kennicutt 1997; Zurita et al. 2002). In the local group, Bland-Hawthorn & Maloney (1999) and Putman et al. (2003) find LyC escape fractions of only 1–2%.

In the hierarchical formation scenario of galaxy assembly, the inflow of cold gas and merging of high-redshift compact galaxies plays a role in the formation of massive young starburst galaxies. The number density of those Luminous Compact Blue Galaxies is also known to increase strongly with redshift (Lilly et al. 1998; Mallén-Ornelas et al. 1999). *FUSE* observations of analogous nearby, young, starbursting dwarf galaxies have shown that the f_{esc} values of these galaxies reach $\lesssim 4\%$, and can collectively contribute a significant fraction to the ionizing background at high redshift (Bergvall et al. 2006; Grimes et al. 2007, 2009; Leitert et al. 2011, 2013). Most theoretical models also predict that low-mass galaxies abundant at high redshifts are more likely to have higher f_{esc} values than the larger galaxies at low to moderate redshift (Razoumov & Sommer-Larsen 2010; Yajima et al. 2011; Wise et al. 2014). Hence, it is likely that different classes of objects dominated reionization at different cosmic epochs, i.e., the combined far-UV output from (dwarf) SFGs may have started to reionize the IGM at $z \gtrsim 6.5\text{--}7$, then, along with more massive galaxies, completed and maintained its ionized state at $z \lesssim 6$ until AGN started to dominate at FUV wavelengths at $z \lesssim 2\text{--}2.5$.

In this work, we describe our analysis of *HST* rest-frame UV observations of LyC escaping from massive SFGs and weak AGN at $z \simeq 2.3\text{--}5.8$ in three UVIS filters with the Wide Field Camera 3 (WFC3) and one optical filter with ACS. We compare the measured LyC fluxes of our sample to their modeled intrinsic LyC fluxes using stellar population synthesis




Figure 1. [a] Example composite rest-frame far-UV spectra of SDSS QSOs at $\langle z \rangle \sim 1.3$ (Vanden Berk et al. 2001 [blue]) and of LBGs at $z \sim 2\text{--}4$ (Bielby et al. 2013 [green and orange]; Shapley et al. 2003 [red]). The WFC3/UVIS F225W, F275W, F336W, and ACS/WFC F435W filter transmission curves are ideally positioned to capture Lyman continuum emission ($\lambda < 912\text{\AA}$) at $z \geq 2.26$, $z \geq 2.47$, $z \geq 3.08$, and $z \geq 4.35$, respectively. The combined SEDs of SDSS QSOs suggest a strong LyC signal, whereas the SEDs of LBGs suggest fainter LyC flux. [b] Total system throughput curves (observed wavelengths) of the same WFC3/UVIS and ACS/WFC filters (Dressel et al. 2015; Avila et al. 2015) are shown on a logarithmic scale to emphasize their out-of-band transmission (“red-leak”). These WFC3 UVIS filters were designed to simultaneously maximize throughput and minimize such red-leaks. In Appendix B.1 we assess the percentage of non-ionizing UVC flux with $\lambda > 912\text{\AA}$ leaking in the filter. We include the ACS/WFC F606W filter as a UVC filter reference.

models and Monte Carlo (MC) simulated line-of-sight IGM transmission models.

This paper is organized as follows. In §2, we describe the data that we used for our analysis and how it was reduced. In §3 we give our assessment of the available spectra for our galaxy samples and their reliability and completeness. In §4, we outline the method we implemented to create the stacked LyC images of our samples of galaxies, how we perform photometry on the stacks, the observed LyC flux that we measure, and the significance of these detections. In §5, we introduce the stacked LyC escape fraction, how we calculated the f_{esc} values, their implications, and the observed and modeled radial profiles of the escaping LyC from our samples. In §6 and §7, we discuss our results and present our conclusions. We use Planck (2015) cosmology throughout: $H_0 = 67.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.308$ and $\Omega_\Lambda = 0.692$. All flux densities (referred to as “fluxes” throughout) quoted are in the AB-magnitude system (Oke & Gunn 1983), unless stated otherwise.

2. WFC3/UVIS AND ACS/WFC OBSERVATIONS AND DATA REDUCTION

2.1. ERS/GOODS-S WFC3/UVIS and ACS/WFC Data

Our data was taken with *HST*’s WFC3/UVIS camera in the Early Release Science (ERS) field in September 2009 (Windhorst et al. 2011, hereafter W11), and in July 2002–May 2003 with ACS/WFC as part of the Great Observatories Origins Deep Survey (GOODS; Dickinson et al. 2003). Fig. 1a and 1b show the WFC3/UVIS F225W, F275W, F336W, and ACS/WFC F435W filter transmission curves, which are ideally positioned to capture LyC emission with negligible red-leak at $z \gtrsim 2.26$, $z \gtrsim 2.47$, $z \gtrsim 3.08$, and $z \gtrsim 4.35$, respectively (see Appendix B.1 for a detailed discussion on red-leak). LyC observations at $z \gtrsim 4$ using optical imaging (e.g.,

ACS/WFC F435W images) are much more difficult to obtain due to the higher IGM opacity at these redshifts. The models of Inoue et al. (2014) suggest that the IGM transmission of ionizing photons at $z \sim 5$ could still reach $\sim 12\%$ in some sight-lines, so we include ACS/WFC F435W imaging of the GOODS-South field (Giavalisco et al. 2004) in our analysis of LyC emission at high redshift as well. The corresponding rest-frame non-ionizing ultraviolet continuum (UVC) imaging of our galaxies were taken with ACS/WFC in the F606W, F606W, F775W, and F850LP filters, respectively. These filters sample rest-frame $\lambda_{\text{eff}} \sim 1400\text{--}1800\text{\AA}$ for each of our redshift intervals. When we model the rest-frame UVC absolute magnitudes (M_{AB}) from the spectral energy distribution (SED) fits, we integrate over the interval $1500 \pm 100\text{\AA}$ (see § 5.1). We also utilized photometry from WFC3/IR F098M, F125W, and F160W imaging in the ERS field (W11) and CANDELS WFC3/IR F105W, F125W, and F160W (Grogin et al. 2011; Koekemoer et al. 2011) photometric catalogs in GOODS-South (Guo et al. 2013) for object selection and SED fitting (see §3.1 and §5.1).

Table 1 summarizes the data in the ERS and GOODS South fields available for studying LyC emission, and the redshift range over which each of these filters can sample LyC emission with negligible contamination from non-ionizing flux. Each lower redshift bound was carefully chosen such that no light with $\lambda > 912\text{\AA}$ is sampled below the filter’s red edge (defined at 0.5% of the filter’s peak transmission, as tabulated in Dressel et al. (2015) and references therein). The upper redshift bound of each bin in Table 1 occurs at the redshift where the next redder filter can trace LyC emission more sensitively. Fig. 1[a] suggests that the observed escaping LyC emission strongly declines towards shorter wavelengths. For this reason, the broadband filters we use are most sensitive to LyC emission at the low-redshift end of each of the four redshift




Figure 2. Residual astrometric errors in the improved WFC3/UVIS ERS mosaics in F225W [left], F275W [middle], and F336W [right], respectively, as measured relative to the ACS/WFC F435W ERS mosaic. The new WFC3/UVIS geometric distortion corrections provided a significantly improved registration of all 8 WFC3/UVIS tiles to the ACS/WFC F435W mosaics compared to the 2009 ERS mosaics of W11. The measured residual systematic offsets are $\langle \Delta X \rangle \lesssim 0''.0022$ and $\langle \Delta Y \rangle \lesssim 0''.0172$ compared to ACS (indicated by the dashed lines), and random deviations less than $0''.054$ in X and $0''.073$ in Y.

ranges of Table 1.

2.2. Image Calibration, Drizzling and Astrometric Accuracy

The photometric and astrometric calibration and drizzling of the ACS/WFC mosaics are described in Giavalisco et al. (2004). The initial astrometric calibration of the WFC3/UVIS ERS mosaics, described in W11, left systematic offsets between individual WFC3 CCDs of up to ~ 5 drizzled pixels ($\sim 0''.5$) compared to the GOODS v2.0¹ F435W mosaics, especially at the edges of each UVIS mosaic tile (see Appendix A of W11). These offsets occurred in part due to the way the ERS UVIS exposures were taken, but were primarily due to the rather uncertain geometric instrument distortion correction (IDC) tables available at the time (Kozhurina-Platais et al. 2009). The lack of UV-bright astrometric reference sources in the shallow (~ 900 – 1400 s) individual exposures further prevented accurate registration and drizzling of the F225W, F275W, and F336W images. This issue was

resolved with the improved IDC tables of Kozhurina-Platais et al. (2013) and Kozhurina-Platais (2014). Using these new IDC tables, we re-drizzled the UVIS ERS images into mosaics at a plate scale of $0''.03 \text{ pix}^{-1}$, which were subsequently rebinned to $0''.09$ pixels, in order to increase our sensitivity to any faint, extended, low surface brightness (SB) LyC signal.

Fig. 2 shows the residual astrometric errors of the new ERS mosaics (which we refer to as ERS “v2.0”) for the F225W, F275W, and F336W filters, measured relative to the ACS/WFC F435W mosaics. Residual systematic offsets from the ACS/WFC F435W GOODS v2.0 mosaics for the 8 WFC3/UVIS ERS tiles are now measured to be $\langle \Delta X \rangle \lesssim 0''.0022$ (0.024 pix) and $\langle \Delta Y \rangle \lesssim 0''.0172$ (0.19 pix), with 1σ random deviations less than $0''.054$ (0.60 pix) in X and $0''.073$ (0.81 pix) in Y. Any remaining systematic astrometric offsets are at the sub-pixel level, and are sufficiently small that they no longer affect our SB sensitivity to LyC flux, nor do they add contamination from neighboring sources that can potentially blend in with the LyC signal due to astrometric uncertainties.

2.3. WFC3/UVIS Residual Sky-Background

The original ground-based WFC3/UVIS thermal vacuum flats left residual gradients and patterns in the sky-background at the 5–10% level (Sabbie 2009; W11). For the reprocessing of the ERS data, we use the flat-fields from Mack et al. (2013), which include on-orbit “delta-flat” corrections to the ground-based thermal vacuum flat-fields, significantly reducing the large-scale flat-field patterns seen in earlier ERS mosaics.

W11 measured zodiacal sky-background SB levels in the ERS of 25.46, 25.64, 24.82, and 23.66 mag arcsec⁻² in the WFC3/UVIS F225W, F275W, and F336W, and ACS/WFC F435W images, respectively. When drizzling the data, this sky-background is subtracted (see Koekemoer et al. 2013). For the 5688, 5688, 2788, and 7200 s total exposure times in these filters, this corresponds to a subtraction of ~ 0.00219 , 0.00202, 0.00704, and 0.0517 $e^-/0''.09$ pixel. To determine the best-fit residual sky background level across the

Table 1. Summary of *HST* WFC3/UVIS and ACS/WFC Images and Image Stacks in the ERS Field

Filter	$\lambda / \Delta\lambda^a$	z_{LyC}^b	Obs. Date	t_{exp}^c	PSF ^d	SB(obs) ^e
F225W	2359/467	2.26–2.47	2009 Sep	5,688	$0''.087$	29.80
F275W	2704/398	2.47–3.08	2009 Sep	5,688	$0''.087$	29.82
F336W	3355/511	3.08–4.35	2009 Sep	2,778	$0''.088$	29.76
F435W	4348/800	4.35–6.0	2002 Aug	7,200	$0''.144$	28.33

^a Central wavelength / bandwidth of filter in Å; ^b Redshift range over which rest-frame LyC emission can in principle be sampled. The high-end of each bin occurs at the redshift where the next redder filter can better sample LyC emission at the same or higher redshift;

^c Average integration time of the mosaics in seconds; ^d Typical stellar PSF FWHM. Note that the PSF is undersampled at the $0''.09/\text{pix}$ plate scale of our mosaics; ^e Measured 1σ surface brightness sensitivity limit of our mosaics for a source of uniform SB in a $2''.00$ diameter aperture in AB-mag arcsec⁻² (see W11).

¹ http://archive.stsci.edu/pub/hlsp/goods/v2/h_goods_v2.0_rdm.html




Figure 3. Residual sky-background levels in the drizzled WFC3/UVIS ERS mosaics in the F225W [left], F275W [middle], and F336W [right] filters, determined as described in §2.3. The best fit to the 2009 ERS mosaics of W11 (“v0.7”) is shown in black, while the improved mosaics discussed here (“v2.0”) are shown in red. Dashed lines show Gaussian fits to the two histograms. Each panel lists the best-fit *residual sky-background level* (in ADU/sec), equivalent to 30.29, 29.99, and 28.15 mag arcsec^{−2}, respectively, and the uncertainties thereon across the entire WFC3 ERS mosaic.

WFC3/UVIS ERS mosaics, we follow Freedman & Diakon (1981), who define the sampled bin width for optimal histogram fitting as $2 \times \text{IQR} \times N^{-1/3}$, where IQR is the inner quartile range of the pixel distribution (i.e., the range within 75% and 25%, or $\pm 1.349\sigma/2$ for a Gaussian distribution), and N is the total number of pixels used in the image to construct that histogram. We fit the logarithm of the sky-pixel histogram between -3σ and $+1\sigma$ to a Gaussian function by least squares to obtain the peak value of the fitted functions.

Fig. 3 shows the sky-pixel histograms and best-fit *residual sky-background* levels in the WFC3/UVIS F225W, F275W, and F336W ERS mosaics of W11 in black, while the red curves and measurements indicate the best fit to the data in the current, improved v2.0 mosaics. The slight narrowing of the negative tail of the Gaussian noise distributions in the new mosaics reflects the better flat-fielding. Our best-fit *residual sky background* values and uncertainties thereon are $(2.62 \pm 0.09) \times 10^{-5}$, $(3.71 \pm 0.10) \times 10^{-5}$, and $(31.94 \pm 0.18) \times 10^{-5} \text{ e}^-/\text{s}$ in the F225W, F275W, and F336W filters, respectively, which corresponds to residual sky SB levels of 30.29, 29.99, and 28.15 mag arcsec^{−2} left in the UVIS images *after* drizzling, which subtracted the sky-background to first order. Compared to the observed ERS sky-backgrounds measured in W11, these residual sky SB level values are 4.84, 4.35, and 3.33 mag fainter than the UV sky (1.2%, 1.8%, 4.7% of the UV sky), respectively. These residual sky-background levels can be accurately determined locally and subsequently subtracted, which we employed in our sub-image stacking technique to further increase our sensitivity to extended, low-SB LyC signal (see §4.1).

3. SPECTROSCOPIC REDSHIFTS AND SAMPLE SELECTION

3.1. Spectroscopic Sample Selection

In order to obtain accurate estimates of LyC escape fractions as low as $f_{\text{esc}} \sim 1.0\%$ or lower, we must require the interloper fraction to be very small. Thus, any potentially contaminating, low redshift, interloping galaxies that might create a false-positive LyC signal must be identified and removed from our sample. We therefore require each galaxy that we include in our analysis to have a highly reliable spectroscopic redshift.

Several wide field ground-based spectroscopic surveys have been performed in the GOODS fields, including the ERS

region, at low and high redshift with the Very Large Telescope (VLT) (e.g., Cristiani et al. 2000; Le Fèvre et al. 2004; Szokoly et al. 2004; Wolf et al. 2004; Vanzella et al. 2008; Popesso et al. 2009; Wuyts et al. 2009; Balestra et al. 2010; Silverman et al. 2010; Fiore et al. 2011; Kurk et al. 2012). We retrieved the reduced 1-dimensional FITS spectral data from the ESO archives² for galaxies at $z \geq 2.26$ in order to verify the designated redshifts of these galaxies. We plotted each spectrum at both the original spectral resolution and smoothed with a $\sigma = 3 \text{ \AA}$ Gaussian convolution kernel to reduce the noise in the spectrum for inspection. Typical spectral emission or absorption features for SFGs and AGN were indicated, and portions of the spectra around these features were magnified for closer examination. Specifically, these emission or absorption features include the Lyman Break at 912 Å, Ly α 1216 Å, Si II 1260 Å, O I 1304 Å, C II 1335 Å, Si IV 1398 Å, C IV 1549 Å, and C III] 1909 Å, and when present, C II] 2326 Å, Fe II 2344 Å, and sometimes N V 1240 Å, Fe II 2600 Å, and Mg II 2798 Å. In addition, we included high-contrast cutout images of the galaxies in the filter sampling the rest-frame LyC emission, and all available longer-wavelength filters for inspection and removal of contaminating objects.

Five of us (BMS, RAW, SHC, RAJ, and LJ) visually inspected all spectra and assigned a fidelity grade of 0 (“Gold”), 1 (“Silver”), or 2 (“Bronze”) to each spectrum, where “Gold” indicates a “high quality” (HQ) spectrum or “very likely correct” redshift, “Silver” indicates an “intermediate quality” (IQ) spectrum or “likely or possibly correct” redshift and “Bronze” indicates an “indeterminable” spectrum or “likely incorrect” redshift. Several objects with significant grade discrepancies were discussed in more detail, resulting in some cases in upward or downward adjustment of individual grader’s assessments. The final quality assessments were then averaged. Fig. 4 shows the final grade distributions of the 5 individual assessors, as well as their average grade distribution. Note that some objects have two or more independent spectra (taken in different slit masks, at different grating settings, or from different spectroscopic surveys), so the numbers in the solid histograms add up to 383 for the initial total of 172

² <http://archive.eso.org/> and <http://www.eso.org/sci/activities/garching/projects/goods/MasterSpectroscopy.html>




Figure 4. Comparison of the independent redshift reliability (spectrum quality) assessments, from best (0.0) to poorest (2.0), by five of the authors [BMS, RAW, SHC, RAJ, and LJ]. The highest fidelity bin (0.00–0.63) are ranked as “Gold” (high quality; HQ), and the next bin (0.64–1.33), with spectroscopic redshifts that are most likely correct ranked “Silver” (intermediate quality; IQ) in a pdf file displaying the spectra of all galaxies. The black dotted histograms in the leftmost five panels represent the average of the final panel; the red dotted histograms represent the average grades for the 172 *unique objects*, rather than for the 252 available *individual spectra*.

galaxies. The redshifts of 64 unique objects were agreed to be of the highest quality (with average grades in the range 0.00–0.63), and compose our primary sample of galaxies and AGN with HQ spectra. Redshifts for another 67 unique objects were ranked as likely correct (with average grades in the range 0.64–1.33), but were not included in our primary analysis, as they did not meet our redshift reliability criteria (See Appendix B.3 for an analysis of the LyC signal from these galaxies combined with our sample of galaxies with HQ spectra).

Note that 14 of the 64 objects with HQ spectra are galaxies hosting a *weak* AGN, as evidenced by Seyfert I or Seyfert II emission lines. We also cross-correlated the positions of our galaxy sample with Chandra 4 Ms and Very Large Array 1.4 GHz source catalogs to identify possible obscured/type II AGN using their radio/X-ray luminosities and photon indices (e.g., Xue et al. 2011; Fiore et al. 2012; Miller et al. 2013; Rangel et al. 2013). We identified two of the 14 galaxies with HQ spectra hosting AGN from their X-ray emission, and three others with IQ spectra. In our analysis, we will consider the subsamples of 50 galaxies without AGN signatures, and 14 galaxies with weak AGN, both separately and combined.

3.2. Completeness and Representativeness of the Spectroscopic Samples

Initially, our sample of galaxies was limited to those with spectroscopic redshifts. Our selection of galaxies with high quality spectra in GOODS-S, which solely determined which objects were included in our analysis, reduced our sample to galaxies that can be observed with ground-based spectroscopy at a high signal-to-noise ratio (SNR). This, of course, can bias our results and their subsequent interpretations, e.g., if f_{esc} is a strong function of luminosity (M_{AB}), dust extinction (A_V), metallicity, and/or age. We therefore must consider how representative the characteristics of our selected galaxy samples are in order to understand differences in the results of our analyses of the populations.

In Fig. 5 we plot the distribution of observed apparent magnitudes (m_{AB}), and the corresponding absolute magnitudes (M_{AB}) of the rest-frame non-ionizing UVC flux ($\lambda_{\text{eff}}=1500\pm100\text{\AA}$) of our samples. We consider all galaxies

(Fig. 5a), galaxies hosting weak AGN (Fig. 5b), and galaxies without AGN (Fig. 5c). These values were derived from the apparent flux of the galaxies at the same rest-frame wavelengths, using their best-fit SED models (see §5.1 and Appendix D.1), so no k-correction is necessary to directly compare the M_{AB} values of the galaxies at various redshifts.

If the spectroscopic samples were complete, their apparent magnitude distributions would resemble the galaxy count distributions of the full V , i , and z -band mosaics (Giavalisco et al. 2004) to a given AB-magnitude limit, since these filters sample the UVC emission indicated in Fig. 5, and because the spectroscopic samples were r -band or i -band selected. Their M_{AB} distribution would also reflect the galaxy UV-luminosity function slope at their effective M_{AB} to the effective completeness limits at these redshifts, which typically sample rest-frame wavelengths $\lambda_{\text{eff}} \sim 1500\text{--}1700\text{\AA}$ (e.g., Reddy & Steidel 2009; Finkelstein et al. 2015).

It is clear from Fig. 5 that our spectroscopically selected samples are incomplete for $m_{\text{AB}} \gtrsim 24.0$ mag, both for galaxies with and without weak AGN. For $m_{\text{AB}} \lesssim 24.0$ mag, the distributions are consistent with the expected slope of the galaxy counts from W11, so the selected samples may be representative for LyC studies, but only for these brighter fluxes. We also note that our selection of galaxies with high SNR spectra will have favored the broad emission lines of (weak) AGN, and Ly α emission or strong absorption line galaxies, while LBGs and other galaxies without prominent spectroscopic features are less likely to have yielded the highly reliable redshifts required to be included in our highest-fidelity sample, even for $m_{\text{AB}} \lesssim 24.0$ mag.

The UVC luminosities of the galaxies in our sample span $-24.2 \lesssim M_{\text{AB}} \lesssim -18.8$ mag, with an average of $M_{\text{AB}} \simeq -20.7^{+0.9}_{-0.7}$ mag (1σ), indicative of predominantly luminous galaxies about as bright as M^* at $2.5 \lesssim z \lesssim 4$ (e.g., Hathi et al. 2010), or of galaxies hosting *weak* AGN. Since this is the only sample for which reliable redshifts currently exist, this is strictly the only luminosity range over which the measurements and analyses of any escaping LyC emission that follows will be valid, which may be more luminous than galaxies that contributed to reionization at $z > 7$ (e.g., Bouwens et al. 2012).

Our sample also does not fully sample the parameter




Figure 5. [a] Absolute and apparent magnitude distributions at the rest-frame $1500 \pm 100 \text{Å}$ of the HQ (red) and HQ+IQ (gray) spectroscopic samples for all galaxies. [b] Same, for just the galaxies without AGN activity. [c] Same, for galaxies with indications of AGN activity. These magnitudes were derived from the observed SED fits (see §5.1), and therefore do not require k-corrections. The blue dashed curve indicates the slope of the luminosity function of $\langle z \rangle = 3.46$ galaxies at $M_{AB} = -20.8$, equal to 0.84 dex/mag.

space of the age, stellar mass, star-formation rate (SFR), and A_V -values for galaxies at these redshifts, indicated by their SED fits (see §5.1). These biases are also more prominent in the individual redshift subsamples. Fig. 6 shows the distribution of these parameters for the HQ spectroscopic sample of galaxies without AGN. The $\langle z \rangle = 2.38$ and $\langle z \rangle = 2.68$ galaxies more or less evenly sample the A_V and age distribution, which ranges from $A_V = 0.0\text{--}2.6$ mag ($\langle A_V \rangle \simeq 0.75$ mag) and $\sim 10^{5.5}\text{--}10^{9.2}$ yr ($\langle t_{\text{age}} \rangle \simeq 10^{8.4}$ yr) respectively, while the $\langle z \rangle = 3.47$ sample is dominated by somewhat older ($\langle t_{\text{age}} \rangle \simeq 10^{8.9}$ yr), lower extinction galaxies ($\langle A_V \rangle \simeq 0.4$ mag), and the $\langle z \rangle = 5.02$ galaxies are generally younger ($\langle t_{\text{age}} \rangle \simeq 10^{7.5}$ yr) and have also lower extinction ($\langle A_V \rangle \simeq 0.4$ mag). We note that the variation in these parameters from sample to sample is most likely due to the selection of the spectroscopic sample, rather than any real correlation in redshift, as the higher redshift galaxies were generally selected in the redder ACS filters. Our sample is also indicative of active star-formation, with SFRs ranging from $\sim 10^{0.8}\text{--}10^{3.6} M_{\odot}/\text{yr}$, while the majority of galaxies are producing stars at $\sim 10^{1.4} M_{\odot}/\text{yr}$. Our galaxies are generally quite massive, with stellar masses ranging from $\sim 10^9\text{--}10^{12} M_{\odot}$ and an average stellar mass of $\sim 10^{10.6} M_{\odot}$, which is also implied by their rather luminous average M_{AB} values. The incomplete sampling of these various parameters should be taken into account when interpreting the f_{esc} values for these individual subsamples. In order to obtain a more representative sample of galaxies in each redshift bin, we must include more galaxies that sample the full range of these parameters at their respective redshifts, with average parameters that reflect the true averages for all galaxies at these redshifts, and probe fainter luminosities. This should be a focus of future deeper spectroscopic surveys, either from the ground or with JWST.

4. LYC EMISSION STACKING AND PHOTOMETRY

4.1. Sub-Image Stacking for each LyC Filter

Since absolute LyC escape fractions have been measured to be very low, and the detected LyC emission to be very faint or not detected at all at $z \lesssim 3$ (e.g., Steidel et al. 2001; Shapley et al. 2006; Iwata et al. 2009; Siana et al. 2010; Mostardi et al. 2013), we apply a technique of sub-image stacking of the *observed* LyC emission from multiple galaxies to increase the total SNR and sensitivity to the faint, potentially low-SB LyC flux from individual galaxies. Stacking LyC emission from galaxies at similar redshifts can be used to quantify the *average* LyC emission from galaxies at their average redshift. This method also reduces small-scale residual systematic errors in the stacked sub-images left from bias, dark current, sky-subtraction, flat-fielding, and/or any gradients from variations in exposure time or photon-noise between exposures that might remain in the background of drizzled mosaics, as effects from random systematics are averaged out in a stack. We create our stacks of the LyC emission for our samples as following.

For each galaxy, we extracted 71×71 -pixel ($6''.39 \times 6''.39$) sub-images from the WFC3/UVIS or ACS/WFC mosaics in the respective filter that samples the LyC emission of each galaxy. The size of these cutouts provided sufficient sampling of the photon statistics in the sub-images for fitting the distribution, while minimizing the potential area of neighboring sources of non-ionizing flux. Each sub-image was centered on the RA and Dec of the flux-weighted centroid of the individual galaxy, as measured in the WFC3/IR F160W ERS mosaic by SExtractor (Bertin & Arnouts 1996).

We then created SExtractor segmentation maps for each LyC sub-image in order to identify *all* objects (including noise peaks) detected at a $\geq 1\sigma$ threshold above the local sky. We then exclude *all* surrounding detections outside of a central circular aperture with a $0''.5$ radius ($r = 5.5$ pix) found in the LyC segmentation maps. We preserve all flux from any objects inside this central aperture when we stack the sub-images, while also excluding those detected on the border of the central aperture. This masking was applied to ensure that all potential sources of non-ionizing flux from lower redshift




Figure 6. Stacked BC03 SED fit parameter distributions of the HQ spectroscopic samples for galaxies without AGN. The purple, blue, green, and red bars represent the $\langle z \rangle = 2.38$, $\langle z \rangle = 2.68$, $\langle z \rangle = 3.49$, $\langle z \rangle = 5.02$ samples, respectively.

neighbors along the line-of-sight are removed before stacking. We note that this procedure would not be possible when stacking LyC emission of galaxies using ground-based observations alone, as effects from seeing can blend neighboring non-ionizing contaminants with the true LyC signal (e.g., Nestor et al. 2013; Siana et al. 2015; Mostardi et al. 2015). Each individual masked sub-image was inspected visually to verify that no surrounding objects indeed remained in the sub-images, including those seen only at longer wavelengths in the 10-band ERS mosaics (see §2). Thus, it is possible that the rigorous removal of surrounding flux can sometimes result in the removal of more extended (i.e., at $r \geq 0''.5$) LyC flux from the stacked images if this were detectable at $\geq 1\sigma$ above the local sky-background (see §4.3 and §5.4).

We also subtracted a constant from each of the sub-images before object removal in order to bring the overall background of the images as close to zero as possible. We calculated this background level from the count-rate histogram of the surrounding pixels outside the central circular aperture, binning their values according to the Freedman-Diaconis rule (see §2.3). We then fit each sky histogram to a Gaussian function by least squares, taking the mean of the fitted Gaussian as the background constant. This local sky-background removal was applied in order to sum the actual LyC flux *above* the background from each sub-image, rather than LyC+background, as variations in background levels between sub-images can suppress the flux contribution from the faintest LyC emission in the stack. The subtraction also removed any residual small scale gradients between the sub-images left from bias/sky-subtraction, flat-fielding, and/or exposure time/noise variations in the mosaics.

We then stacked the processed sub-images of all the galaxies in each UVIS/ACS image with spectroscopic redshifts, where LyC can be observed in their respective filter, by *averaging* all of the pixel count rates of the sub-images. We then created stacked weight maps for each LyC stack by summing the inverse of the pixel values of the corresponding region in the ASTRODRIZZLE (Fruchter et al. 2010; Gonzaga et al. 2012) weight maps where the galaxy sub-images were extracted, then inverting the sum to generate the stacked weight maps (i.e., $W_j = 1/\sum_i 1/W_{i,j}$, where W_j is the weight for

pixel j in sub-image i). These weight maps give the relative weight of each pixel in the LyC stacks, and are used only for quantifying all photometric errors in the observations. We also created median combined stacks by calculating the median value of each pixel in all the processed sub-images, as well as weighted average stacks by summing the weighted pixel values of the processed sub-images, normalized by the sum of their weights (i.e., $\langle f_j \rangle = \sum_i W_i f_{i,j} / \sum_i W_i$, where f_j

represents the flux in counts per second measured in pixel j for sub-image i). We found that the stacks created by averaging without weighting provided the highest SNR and best SB-limit compared to the median and weighted average stacks, as these latter stacks do not include all the available LyC flux from the sub-images. The average stacks also simplified the error analysis for the photometry of the stacks, as well as the subsequent interpretation of the measured LyC signals. We created stacks for the total sample of galaxies and separate stacks for the galaxy and AGN samples, since each sample likely produces the majority of their LyC photons by different mechanisms, which must be taken into account when determining f_{esc} for these galaxies.

Since many f_{esc} values quoted in the literature are calculated relative to the rest-frame non-ionizing UV-continuum (UVC) flux measured from $1500 \lesssim \lambda_0 \lesssim 1700 \text{ \AA}$ (see §5.1), we created corresponding UVC stacks for each LyC stack from sub-images extracted from the ACS/WFC mosaics of the ERS/GOODS-S fields that sample the UVC emission of our galaxies. For the redshift intervals that sampled LyC emission in the F225W, F275W, F336W, and F435W stacks, the ACS/WFC filters that sample the UVC emission correspond to F606W, F606W, F775W, and F850LP, respectively.

The galaxies stacked in the WFC3/UVIS F225W filter contain co-added sub-images frames of 14 galaxies over the redshift-range $2.302 \leq z \leq 2.450$ ($\langle z \rangle = 2.380$), the F275W stack contains 11 galaxies at $2.559 \leq z \leq 3.076$ ($\langle z \rangle = 2.682$), the F336W stack contains 11 co-added galaxies at $3.132 \leq z \leq 3.917$ ($\langle z \rangle = 3.494$), and the ACS/WFC F435W stack contains 15 galaxies at $4.414 \leq z \leq 5.786$ ($\langle z \rangle = 5.015$). These stacks, as well as the corresponding UVC stacks, are shown in Fig. 7. Stacks for the subsamples of galaxies with weak AGN and galaxies without AGN are shown in Fig. 8 and




Figure 7. Sub-image stacks for the four different redshift bins in our sample of *all* galaxies, sampling LyC emission in: [a] F225W at $2.26 \leq z \leq 2.47$, [b] F275W at $2.47 \leq z \leq 3.08$, [c] F336W at $3.08 \leq z \leq 4.35$, and [d] F435W at $4.35 \leq z \leq 6$; and corresponding UVC ($\sim 1400 \lesssim \lambda_0 \lesssim 1800 \text{\AA}$) emission in: [e] F606W, [f] F606W, [g] F775W, and [h] F850LP. Note that the objects contributing to panels [e] and [f] differ, since they correspond to different redshift bins. Blue ellipses indicate the SEXTRACTOR LyC apertures for a 1σ detection criterion, while green ellipses correspond to the SEXTRACTOR MAG_AUTO apertures, based on the centroids and apertures defined in the UVC stacks. All sub-images are 71×71 pixels ($6''.39 \times 6''.39$) in size.




Figure 8. As Fig. 7 for galaxies *without* AGN (i.e., no obvious signs of nuclear activity from their spectra or X-ray/radio luminosities/photon indices).




Figure 9. As Fig. 7 for only the galaxies hosting weak AGN.

9, with elliptical apertures indicating regions where photometry was performed.

The deepest galaxy counts in J and H -band of W11 give us an estimate the total number of contaminating objects that could be present in our $\sim 0''.5$ radius LyC apertures (Fig. 7–9). To the ERS limit of J , $H \lesssim 27.55$ – 27.25 mag, respectively, there are $\lesssim 5.2 \times 10^5$ galaxies deg^{-2} , yielding a $\lesssim 3\%$ probability of finding one unrelated foreground object in, or overlapping with, the LyC aperture. For our sample of 50 galaxies, this would amount to $\lesssim 2$ interlopers. Due to the possible interloper’s proximity to the LyC candidate, it is not always possible to obtain reliable spectroscopic or photometric redshifts for these neighbors. Nonetheless, in these few cases, light from the nearby neighbors was masked out with SEXTRACTOR segmentation maps and repeated for *all* other objects in the 71×71 pixel image sections outside the central $\sim 0''.5$ aperture, to exclude contaminating objects in the photometry in the central aperture, and to assure that accurate measurements of the surrounding sky could always be obtained.

4.2. Rest-frame Lyman Continuum Photometry

The results from our photometry measured in the apertures shown in Fig. 7–9 are summarized in Table 2. Because the LyC flux escaping from galaxies in these stacks is very faint, we perform all of our photometry on the LyC stacks with SEXTRACTOR using detection images in dual-image mode. As our reference images, we use the corresponding non-ionizing UVC stacks and the LyC stacks convolved with a 1σ Gaussian kernel to increase detection of the LyC faintest emission. The *un-convolved* LyC images were used for measurement of LyC fluxes in both cases. The (green) UVC apertures were used to determine the corresponding LyC emission relative to the UVC flux and minimize the inclusion of noise in the measurement. However, since the LyC flux is typically more extended than the UVC (see Fig. 11), we also measure the LyC flux by independently running detection on the smoothed LyC image in order to maximize the detection of LyC flux, while minimizing the associated noise in the measurement. Thus, while the UVC apertures may include more background noise in the measurement, they also do include more of the available LyC signal in the stack. The (blue) LyC apertures attempt to capture most of the LyC flux in the stack, but do include a minimized amount of background noise. Therefore, the optimized LyC apertures typically provide higher SNR than the UVC apertures.

We used the relative weights from the stacked weight-maps for assessing photometric uncertainties. This approach allows us to directly estimate the measurement error from the LyC stack itself, rather than using the absolute variance per pixel in the stack, which can increase substantially for a stack composed of different regions of an image taken with different areas of the detector. We measure the flux from objects in the stacks detected at the $>1\sigma$ level in 5 contiguous pixels above the local sky-background, determined by median filtering the image with a 3×3 pixel kernel in a 32 pixel mesh. We convert the flux measured by SEXTRACTOR to AB magnitudes, using the infinite aperture zeropoints listed on the STScI instrument

websites³.

From the MAG_AUTO parameter measured by SEXTRACTOR, we detect average LyC flux from galaxies and AGN at $m_{\text{AB}} \simeq 29.92, 29.61, 30.13$, and 29.51 mag for the F225W, F275W, F336W, and F435W stacks with measured SNR values at $\sim 6.53, 10.4, 4.56$, and 4.87 in the (blue) LyC apertures indicated in Fig. 7, respectively. We find similar LyC flux in the (green) UVC apertures at $m_{\text{AB}} \simeq 30.01, 29.31, 29.82$, and 30.70 mag (with measured SNR $\sim 4.93, 10.3, 6.01$, and 2.25), respectively. For only the galaxies without AGN (Fig. 8), we measure $m_{\text{AB}} \simeq 29.98, 30.09, 30.79$, and 30.37 at SNR $\sim 5.64, 5.71, 3.91$, and 3.28 , while the flux from galaxies with AGN was measured at $m_{\text{AB}} \simeq 30.12, 28.92, 29.53$, and 28.58 , with SNR $\sim 2.34, 8.77, 5.17$, and 4.48 , respectively (Fig. 9). For stacks with comparable SNR, the AGN stacks are seen to be brighter than the galaxies by $m_{\text{AB}} \sim 1$ mag on average, despite having fewer contributing sub-images in the stacks.

For the stacks of all galaxies, the difference between the LyC flux measured in the LyC-defined and the UVC-defined apertures is $\Delta m_{\text{LyC}}(\text{LyC-UVC apertures}) \simeq +0.05 \pm 0.11$ mag (m.e.) on average. Hence, both apertures measure consistent LyC fluxes to within their respective errors. This effect can be attributed to the spatial distribution of LyC flux in the stacks (see §5.4). We note that, although real LyC flux might exist at the $\sim 1\sigma$ level outside the measurement apertures, we do not incorporate this flux into our measurement, as this would require us to increase our aperture size and add extra noise in the aperture, which would increase the uncertainty of our measurements, as well as the interloper contribution.

We performed a series of critical tests on our data to ascertain the robustness and validity of our stacking procedures and LyC detections, for which we refer the interested reader to Appendix A. From these tests, we conclude that our measurements are reliable to within their measured errors, and are not the result of various possible sources of spurious signal.

4.3. Super-Stacks of LyC Emission from Galaxies at $2.3 \leq z \leq 5.8$

In order to determine at what SNR our observations can measure the LyC flux from our total sample of galaxies that span the $2.3 \leq z \leq 5.8$ redshift range, and probe the faintest LyC emission from our galaxies, we construct a stack of the observed LyC flux in f_ν of *all* the galaxies in our sample as follows. We first extract the sub-images from the F225W, F275W, F336W, and F435W mosaics and apply our sky-subtraction and neighbor masking procedures, as outlined in §4.1. We then convert the pixel count rates of each sub-image from counts/second (cps) to $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$ by scaling the pixel values by their AB magnitude zeropoint, so that all sub-images are weighted by their relative observed fluxes. We then stack the sub-images by averaging, as described in §4.1. We created stacks for the full sample of all galaxies and separate stacks for the galaxies with and without AGN, both for our HQ and HQ+IQ spectra samples, as shown in Fig. 10. We note that these super-stacks represent the average *observed* LyC flux from our sample integrated from $z = 2.3$ – 5.8 through the various sight-lines, and so the flux in these stacks will be

³ http://www.stsci.edu/hst/wfc3/phot_zp_lbn

Table 2. LyC Stack Photometry

Filter	z -range	$\langle z \rangle$	N_{obj}	LyC apertures				UVC apertures				
				m_{LyC}	$\text{ABerr}_{\text{LyC}}$	SNR_{LyC}	A_{LyC}	m_{LyC}	SNR_{LyC}	A_{UVC}	m_{UVC}	SNR_{UVC}
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
ALL GALAXIES:												
F225W	2.291–2.450	2.374	15	29.92	0.17	6.53	0.958	30.01	4.93	1.407	24.50	240.8
F275W	2.470–3.076	2.688	18	29.61	0.10	10.4	0.782	29.31	10.3	1.427	24.68	226.2
F336W	3.132–3.917	3.446	14	30.13	0.24	4.56	0.943	29.82	6.01	0.923	24.75	131.0
F435W	4.414–5.786	4.989	17	29.51	0.22	4.87	0.874	30.70	2.25	0.468	25.79	95.3
GALAXIES WITHOUT AGN:												
F225W	2.302–2.450	2.380	14	29.98	0.19	5.64	1.059	30.00	4.80	1.451	24.43	237.5
F275W	2.559–3.076	2.682	11	30.09	0.19	5.71	0.656	29.80	4.90	1.583	24.51	192.2
F336W	3.132–3.917	3.494	10	30.79	0.28	3.91	0.237	30.64	2.87	0.622	24.96	111.6
F435W	4.414–5.786	5.015	15	30.37	0.33	3.28	0.354	30.61	2.32	0.467	26.12	70.3
GALAXIES WITH AGN:												
F225W	2.291–2.291	2.291	1	30.12	0.46	2.34	0.213	30.00	1.10	1.424	27.90	7.85
F275W	2.470–3.008	2.697	7	28.92	0.12	8.77	1.372	29.56	6.97	0.665	25.00	156.9
F336W	3.217–3.474	3.326	4	29.53	0.21	5.17	0.612	29.34	5.47	0.777	24.52	102.9
F435W	4.760–4.823	4.792	2	28.58	0.24	4.48	0.571	>31.5	<2	0.357	24.66	79.0

Table columns: (1): WFC3 or ACS filter used; (2): Redshift range of galaxies included in LyC/UVC stacks; (3): Average redshift of stack; (4): Number of galaxies with high quality spectroscopic redshifts used in the stacks; (5) Observed average AB magnitude of LyC emission from stack (using the SEXTRACTOR MAG_AUTO aperture, as indicated by the blue ellipses in Figs. 7–9; (6): 1σ error of average LyC AB-mag; (7): Measured SNR of the LyC stack flux (MAG_AUTO); (8): Area (in arcsec²) of the LyC MAG_AUTO (blue) aperture (see Figs. 7–9); (9): Observed total AB magnitude of LyC emission from stack (SEXTRACTOR aperture matched to UVC, indicated by the green ellipses in Figs. 7–9; (10): Measured SNR of the LyC stack flux (matched UVC aperture); (11): Area (in arcsec²) of the UVC aperture; (12): Observed total AB magnitude of the UVC stack (SEXTRACTOR MAG_AUTO aperture); (13): Measured SNR of the UVC stack (SEXTRACTOR MAG_AUTO aperture).

dominated by the galaxies with the brightest apparent LyC flux.

Due to the very low sky-background in the F225W and F275W filters (W11), the relative scaling of the count rates in the sub-images slightly amplified the contribution of detector-noise from these filters in the stack. However, since the F336W and F435W filters are dominated by photon-noise from the much brighter zodiacal background at these wavelengths, the stacks are therefore dominated by sky-background as well. We find that the stack of the total HQ sample reaches a SNR of 11.1σ , while the LyC stacks of HQ galaxies with and without AGN achieve SNR ratios of 11.4σ and 6.8σ in the optimized apertures, respectively. We also find that the total apparent AGN LyC flux outshines that from galaxies without AGN by a factor of $F_{\nu}^{\text{AGN}}/F_{\nu}^{\text{Gal}} \simeq 3.0$ ($\Delta m_{\text{AB}} \simeq 1.2$ mag).

Since these super-stacks were created solely for the purposes of visualization and probing the SNR of the total observed LyC signal from our samples, we do not perform any further analysis of the LyC flux measured in these stacks. Because the absorption of ionizing photons by the IGM is non-linearly dependent on redshift, modeling of the combined intrinsic LyC flux from galaxies spanning $2.3 \lesssim z \lesssim 5.8$ through various sight-lines would become increasingly divergent, such that the resulting total f_{esc} value of these galaxies would be highly uncertain. We therefore only use these super-stacks for total observed flux and SNR comparisons between subsamples and for visualization of the LyC signal from these galaxies.

This exceedingly faint LyC emission emerging from the average of the 50 galaxies without AGN appears to have a flat spatial distribution that is *not* centrally concentrated. This may allude to the manner in which LyC escapes from galaxies. In order to ionize the IGM, LyC photons must escape through holes in the surrounding gaseous and/or dusty material between stars and the IGM, which can be distributed randomly within galaxies. With at most a few clear sight-lines per galaxy, these stacked images suggest that the most likely escape paths of LyC may be on average somewhat offset from the galaxy center, i.e., escaping more from the outskirts than the centers of these galaxies. Given the random orientation of galaxies in each stack, this would explain the faint, non-centrally concentrated, and extended morphology of the detected LyC emission. The LyC stack of all AGN shows a more asymmetric flux distribution compared to the stack of all galaxies without AGN. This may indicate that LyC photons created by accretion disks in AGN escape from galaxies with weak AGN in a preferred direction. Note that the AGN “super-stack” is dominated by flux from the two $z > 4.7$ AGN. We discuss the radial profiles of galaxies further in §5.4.

In our adopted Planck cosmology, the angular size scale decreases by $\sim 23\%$ and the apparent fluxes dim by $\sim 71\%$ from $z \simeq 2.3$ to $z \simeq 5.0$. We deliberately did *not* scale any of the pixel values or resample the pixel-scale to account for these changes during the stacking process over all redshifts, as we only created these super-stacks to quantify the SNR of the average *observed* LyC flux for our entire sample. Using a varying pixel scale for each galaxy would have introduced




Figure 10. [Top Row]: Galaxies in our sample with high quality spectra and reliable redshifts; [Bottom Row]: Galaxies in our sample with both high quality and intermediate quality spectra, containing predominantly reliable, and some possibly incorrect redshifts; *a/e* [Left column of panels:] Stacked LyC emission of all galaxies with spectroscopic redshifts in the ERS field; *b/f* [2nd column:] Stacked LyC emission of all galaxies with (weak) AGN; *c/g* [3rd column:] Stacked LyC emission of all galaxies with no indication of AGN activity; *d/h* [4th column:] Stacks from the 3rd column smoothed with a 1σ Gaussian kernel. These stacks represent the average *observed* LyC F_ν from all galaxies integrated from $2.3 \lesssim z \lesssim 5.8$. Thus, we did not scale the flux from each galaxy to have the same luminosity distance, nor did we resample each sub-image to have the same angular size. The blue circles indicate the region where flux was preserved while stacking, and the green circles show the excluded region where the sky was computed and subtracted ($r=0''.5$ and $1''.08$, respectively), and are centered on the light centroid measured in F125W. The measured SNR of the combined LyC emission in these stacks is $\sim 11.4\sigma$ for the stack of all galaxies with HQ spectra, and $\sim 6.8\sigma$ for the stack containing both high and intermediate quality spectra. The AGN stacks exhibit both a centrally concentrated and extended component in their flux distributions, from contributions of a central AGN point source and perhaps also from scattered photons. The fainter LyC emission emerging from the average of the 50 (114) galaxies without AGN activity appears to display a flat light profile with increasing radius. Hence, LyC emission at $z \simeq 2.3\text{--}5.8$ may *on average* escape along a few random sight-lines through a porous ISM, and need not be centrally concentrated, having at most a few clear sight-lines per galaxy. These images suggest that the likeliest LyC escape paths may be slightly offset from a galaxy center. Given the random orientation of galaxies in each stack, this would explain the faint, non-centrally concentrated, and extended morphology of the detected LyC emission.

correlated inter-pixel resampling noise, which would also decrease the sky SB limits and the effect of averaging over residual subtle systematics. Stacking with the same plate scale for all redshifts also preserves the observed photon statistics, which are needed for accurate sky-subtraction. Hence, resampling all the images as a function of redshift would reduce the SNR of the resulting stacked LyC signal. We note that the physical scale of the galaxies that we stack changes by $\pm 4\%$ within each redshift bin and by $\pm 23\%$ for the entire sample. This does not noticeably affect the LyC and UVC light-profiles in each of our four redshift bins, which are discussed in §5.4, but does “blur” the light distribution seen in the super-stacks in Fig. 10 by approximately these amounts in the radial direction from the center. Fig. 10 can thus only be used to visualize the combined *observed* LyC signal over the entire redshift range $z \simeq 2.3\text{--}5.8$, but cannot be used for further quantitative analysis.

5. LYC ESCAPE FRACTIONS, AND RADIAL SURFACE BRIGHTNESS PROFILES

5.1. Relative and Absolute LyC Escape Fractions for Stacks

Estimating the escape fraction of LyC photons from galaxies, f_{esc} , is non-trivial, as it requires modeling of their apparent *intrinsic* LyC flux, $F_{\nu, \text{LyC}}^{\text{int}}$, and the wavelength-dependent transmission of LyC photons through the IGM, $\mathcal{T}_{\text{IGM}}^{\text{LyC}}(z, \nu)$, for a galaxy at redshift z . Quantitatively, the average *observed* LyC flux ($F_{\nu, \text{LyC}}^{\text{obs}}$) from a stack of galaxies, measured by a photon counting device such as a CCD, is given by:

$$\langle F_{\nu, \text{LyC}}^{\text{obs}} \rangle = \frac{1}{N_{\text{gal}}} \sum_{i=1}^{N_{\text{gal}}} \frac{\int T_{\text{obs}}^{\text{LyC}}(\nu) \mathcal{T}_{\text{IGM}}^{\text{LyC}}(z_i, \nu) f_{\text{esc}, i}^{\text{abs}} F_{\nu, i}^{\text{int}}(\nu) \frac{d\nu}{\nu}}{\int T_{\text{obs}}^{\text{LyC}}(\nu) \frac{d\nu}{\nu}} \quad (1)$$

where N_{gal} denotes the number of galaxies in the stack, $F_{\nu, i}^{\text{int}}$ denotes the *intrinsic* — i.e. produced — stellar SED from galaxy ‘i’, $f_{\text{esc}, i}^{\text{abs}}$ denotes the fraction of the observed LyC flux that escaped from the galaxy into the IGM, $\mathcal{T}_{\text{IGM}}^{\text{LyC}}(z_i, \nu)$ denotes the wavelength-dependent IGM transmission curve for galaxy ‘i’ at redshift z , which we acquired from the recent absorption models of Inoue et al. (2014), and the $T_{\text{obs}}^{\text{LyC}}(\nu)$ term denotes the combined transmission of the throughput from the Optical Telescope Assembly (OTA), the filter throughput, and the Quantum Efficiency (QE) of the detector used for the LyC ob-

servation (see Fig. 1 for the WFC3/UVIS and ACS/WFC total system throughput curves).

We expect f_{esc} generally to vary between individual objects. However, since we stack the observed LyC emission from all galaxies in our sample, we simplify our analysis by assuming a constant $f_{\text{esc}}^{\text{abs}}$ value for all galaxies within a given redshift bin. We denote this ‘sample averaged’ escape fraction with $\langle f_{\text{esc}}^{\text{abs}} \rangle$. We can then take $\langle f_{\text{esc}}^{\text{abs}} \rangle$ outside of the sum in Eq 1, and write:

$$\langle f_{\text{esc}}^{\text{abs}} \rangle = \frac{\langle F_{\nu, \text{LyC}}^{\text{obs}} \rangle}{\frac{1}{N_{\text{gal}}} \sum_{i=1}^{N_{\text{gal}}} \frac{\int T_{\text{obs}}^{\text{LyC}}(\nu) \mathcal{T}_{\text{IGM}}(z_i, \nu) F_{\nu, i}^{\text{int}}(\nu) \frac{d\nu}{\nu}}{\int T_{\text{obs}}^{\text{LyC}}(\nu) \frac{d\nu}{\nu}}}, \quad (2a)$$

which can also be expressed as:

$$\langle f_{\text{esc}}^{\text{abs}} \rangle = \frac{\langle F_{\nu, \text{LyC}}^{\text{obs}} \rangle}{\frac{1}{N_{\text{gal}}} \sum_{i=1}^{N_{\text{gal}}} F_{\nu, \text{LyC}, i}^{\text{int}}} = \frac{\langle F_{\nu, \text{LyC}}^{\text{obs}} \rangle}{\langle F_{\nu, \text{LyC}}^{\text{int}} \rangle}. \quad (2b)$$

A more thorough analysis should also take into account that the impact of the IGM varies substantially around this mean for individual objects and sight-lines (see Inoue & Iwata 2008; Nestor et al. 2011). We determine $F_{\nu, i}^{\text{int}}(\nu)$ for each galaxy from the minimized χ^2 fit Bruzual & Charlot (2003, BC03) synthetic stellar population model SED, which was fit to 4–6 non-ionizing continuum broadband WFC3/IR+ACS/WFC measurements at the known *fixed* redshift. These best-fit SEDs were allowed four degrees of freedom for the χ^2 minimization at the fixed spectroscopic redshift of each object, i.e., the age, stellar mass, A_V , and the exponentially decreasing SFR timescale (τ). These were fit from a grid of SEDs using a wide range in each of these parameters. Thus, the best-fit SEDs correspond to the observed F_{ν} of the galaxy with Calzetti et al. (2000) attenuation applied to the *intrinsic* SED, thereby determining the best-fit A_V -value. We use a Salpeter (1955) IMF and adopt solar metallicities for the SEDs.

In order to determine the absolute fraction of escaping LyC ($f_{\text{esc}}^{\text{abs}}$), which compares the apparent flux of LyC photons produced by stars in the galaxy to the observed LyC emission (i.e., $F_{\nu, \text{LyC}}^{\text{obs}}/F_{\nu, \text{LyC}}^{\text{int}}$), the effects from dust must be removed from the SED to obtain its *intrinsic* LyC flux, $F_{\nu, \text{LyC}, i}^{\text{int}}(\nu)$. Since we began our SED fitting with the intrinsic stellar population photospheric flux — which was then reddened by a specified A_V -value using Calzetti et al. (2000) attenuation — we simply use the initial dust-free intrinsic stellar photospheric SED to calculate $F_{\nu, i}^{\text{int}}(\nu)$ for each galaxy. Thus, the *absolute* escape fraction quantifies the amount of LyC that is not absorbed by dust, the multiphase ISM, or other sources of LyC absorption in the galaxy. The fraction of escaping LyC photons that are absorbed by components of the ISM aside from dust, $\langle f_{\text{esc}}^{\text{abs}} \rangle_{\text{red}}$, can also be determined by including the modeled dust attenuation of the SED in Eq. 2a.

We can also determine the escape fraction of LyC flux *relative* to the non-ionizing UVC flux ($F_{\nu, \text{UVC}}$), denoted as $f_{\text{esc}}^{\text{rel}}$. We define the sample average relative LyC escape fraction of

a stack as:

$$\langle f_{\text{esc}}^{\text{rel}} \rangle = \frac{1}{N_{\text{gal}}} \sum_{i=1}^{N_{\text{gal}}} \frac{F_{\nu, \text{UVC}, i}^{\text{int}}/F_{\nu, \text{LyC}, i}^{\text{int}}}{F_{\nu, \text{UVC}, i}^{\text{obs}}/F_{\nu, \text{LyC}, i}^{\text{obs}}} \simeq \frac{1}{N_{\text{gal}}} \frac{\langle F_{\nu, \text{LyC}}^{\text{obs}} \rangle}{\langle F_{\nu, \text{UVC}}^{\text{obs}} \rangle} \sum_{i=1}^{N_{\text{gal}}} \frac{F_{\nu, \text{UVC}, i}^{\text{int}}}{F_{\nu, \text{LyC}, i}^{\text{int}}} \quad (3a)$$

Using Eq. 2b, we can further simplify this expression as:

$$\langle f_{\text{esc}}^{\text{rel}} \rangle \simeq \langle f_{\text{esc}}^{\text{abs}} \rangle \frac{\sum_{i=1}^{N_{\text{gal}}} F_{\nu, \text{UVC}, i}^{\text{int}}}{\sum_{i=1}^{N_{\text{gal}}} F_{\nu, \text{UVC}, i}^{\text{obs}}} = \langle f_{\text{esc}}^{\text{abs}} \rangle \left\langle \frac{F_{\nu, \text{UVC}}^{\text{int}}}{F_{\nu, \text{UVC}}^{\text{obs}}} \right\rangle, \quad (3b)$$

where $F_{\nu, \text{UVC}, i}^{\text{obs}}$ is the observed UVC flux from galaxy ‘i’ as measured in the ACS/WFC UVC filters (see §4.1) and:

$$F_{\nu, \text{UVC}, i}^{\text{int}} = \frac{\int T_{\text{obs}}^{\text{UVC}}(\nu) F_{\nu, i}^{\text{int}}(\nu) \frac{d\nu}{\nu}}{\int T_{\text{obs}}^{\text{UVC}}(\nu) \frac{d\nu}{\nu}} \quad (4)$$

for UVC observations with a total system throughput of $T_{\text{obs}}^{\text{UVC}}(\nu)$. Thus, the relative and absolute escape fractions differ by a factor of $\left\langle \frac{F_{\nu, \text{UVC}}^{\text{int}}}{F_{\nu, \text{UVC}}^{\text{obs}}} \right\rangle = \langle f_{\text{esc}}^{\text{UVC}} \rangle^{-1}$ for the total sample, which deviates from unity depending on the A_V and χ^2 values of the SED fits. The escape fraction of non-ionizing UVC photons, $f_{\text{esc}, i}^{\text{UVC}}$, is related to the observed reddening in galaxy ‘i’ as $f_{\text{esc}}^{\text{UVC}} = 10^{-0.4 A_{\text{UVC}}}$. Note that this term can be omitted from Eq. 3a when using the *intrinsic* (unreddened) model SEDs instead of the observed ones. For a simple “direct” calculation of $f_{\text{esc}}^{\text{rel}}$ that can be computed from values listed in Table 3, using Eq. 3a, we can take the stacked observed and intrinsic LyC and UVC flux ratios of the galaxies, along with the average IGM LyC transmission values, as:

$$\langle f_{\text{esc}}^{\text{rel}} \rangle_{\text{dir}} = \frac{\langle \frac{F_{\text{UVC}}}{F_{\text{LyC}}} \rangle_{\text{int}}}{\langle \frac{F_{\text{UVC}}}{F_{\text{LyC}}} \rangle_{\text{obs}}} \langle \mathcal{T}_{\text{IGM}}^{\text{LyC}} \rangle^{-1}. \quad (5)$$

We determine the ratio of intrinsic fluxes of the LyC and UVC emission from all galaxies as:

$$\left\langle \frac{F_{\text{UVC}}}{F_{\text{LyC}}} \right\rangle_{\text{int}} = \frac{\sum_{i=1}^{N_{\text{gal}}} F_{\nu, \text{UVC}, i}^{\text{int}}}{\sum_{i=1}^{N_{\text{gal}}} F_{\nu, \text{LyC}, i}^{\text{int}}} \quad (6)$$

without applying the $\mathcal{T}_{\text{IGM}}(z_i, \nu)$ term to $F_{\nu, \text{LyC}, i}^{\text{int}}$. The *observed* LyC and UVC flux ratios can be obtained by performing photometry on stacked images of the LyC and UVC emission shown in figures 7–9.

As a specific example of the “direct” calculation of the relative escape fractions, let us consider the F275W filter measurements for the redshift interval with $\langle z \rangle = 2.68$. We use the LyC aperture (blue ellipses in Fig. 7–9), since they generally have the highest SNR (see Table 2). For the stack of all galaxies without AGN, we measure a formal SNR of 5.7σ for a LyC flux of $m_{\text{AB}}^{\text{LyC}} = 30.09$ mag in

Table 3. Summary of f_{esc} Constraints

$\langle z \rangle$	N_{obj}	$\langle f_{\text{UVC}}/f_{\text{LyC}} \rangle_{\text{obs}}$	$\langle f_{\text{UVC}}/f_{\text{LyC}} \rangle_{\text{int}}$	$\langle t_{\text{age}} \rangle$	$A_{V \text{ med}}$	$\langle \mathcal{T}_{\text{IGM}} \rangle$	$\langle f_{\text{esc}}^{\text{rel}} \rangle_{\text{dir}}$	$\langle f_{\text{esc}}^{\text{rel}} \rangle$	$\langle f_{\text{esc}}^{\text{abs}} \rangle_{\text{red}}$	$\langle f_{\text{esc}}^{\text{abs}} \rangle$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
GALAXIES WITHOUT AGN:										
2.380	14	157_{-16}^{+44}	$3.45_{-0.11}^{+0.12}$	$10_{-0.4}^{+0.8}$	$0.55_{-0.44}^{+0.70}$	$0.297_{-0.083}^{+0.082}$	$6.3_{-1.2}^{+3.8}$	$3.5_{-2.1}^{+67.5}$	$30.5_{-12.9}^{+39.1}$	$0.10_{-0.06}^{+1.97}$
2.682	11	162_{-17}^{+45}	$2.98_{-0.07}^{+0.08}$	$10_{-0.7}^{+0.7}$	$0.58_{-0.40}^{+0.89}$	$0.247_{-0.085}^{+0.086}$	$6.0_{-1.1}^{+5.2}$	$3.1_{-1.6}^{+32.4}$	$58.6_{-26.6}^{+77.5}$	$0.14_{-0.07}^{+1.46}$
3.472	11	190_{-21}^{+76}	$11.4_{-0.2}^{+0.2}$	$10_{-0.4}^{+0.7}$	$0.18_{-0.12}^{+0.64}$	$0.112_{-0.050}^{+0.050}$	30_{-16}^{+54}	$19.8_{-10.6}^{+39.2}$	$21.4_{-11.4}^{+46.1}$	$17.6_{-9.8}^{+31.3}$
5.015	15	$43.1_{-4.7}^{+29.2}$	$3.59_{-0.29}^{+0.34}$	$10_{-0.7}^{+0.5}$	$0.17_{-0.12}^{+0.67}$	$0.00108_{-0.00107}^{+0.00123}$	>100	>100	>100	$4.0_{-4.0}^{+96.0}$

Table columns: (1): Average redshift of each stack; (2): Number of objects in each redshift bin, as in Table 2; (3): Average *observed* flux ratio $f_{\text{UVC}}/f_{\text{LyC}}$ and its $\pm 1\sigma$ error range, as measured from the LyC and UVC stacks in their respective apertures (see §4.2 and Table 2); (4): Average *intrinsic* flux ratio $f_{\text{UVC}}/f_{\text{LyC}}$ and its $\pm 1\sigma$ error range, as derived from the BC03 best-fit SED models galaxies without AGN in each of our redshift bins (see §5.1 and Eq. 6); (5): Median dust extinction A_V and its $\pm 1\sigma$ error range of the best-fit BC03 SED model (the median A_V is more representative, as the distributions of each subsample is asymmetric; see §D.1); (6): Average age of the stellar populations from the best-fit BC03 models and their $\pm 1\sigma$ standard deviations in years; (7): Average filter-weighted IGM transmission of all sight-lines and redshifts in the stacks and their $\pm 1\sigma$ standard deviations, calculated from the Inoue et al. (2014) models; (8) Inferred *relative* escape fraction of LyC photons in percent plus and its $\pm 1\sigma$ error range, assuming a constant *intrinsic* LyC to UVC flux ratio and LyC IGM transmission, as described in §5.1 (Eq. 5). The superscript “rel” indicates that the escape fractions were calculated relative to the escape of non-ionizing UVC (eq. 3b). The dust-corrected (intrinsic) stellar SEDs were used to calculate $\langle f_{\text{UVC}}/f_{\text{LyC}} \rangle_{\text{int}}$, so no further dust-corrections were needed, although uncertainties in the SED were included in these values. The stellar SED fits of §5.1 used in Cols. (4)–(6) only include Galaxies *without* weak AGN. (9) The resulting ML and $\pm 1\sigma$ values of the Monte Carlo analysis on $f_{\text{esc}}^{\text{rel}}$ in percent, as described in §5.1 (Eq. 3a); (10) ML and $\pm 1\sigma$ values of the Monte Carlo analysis of $f_{\text{esc}}^{\text{abs}}$ with dust attenuation applied, i.e., the escape fraction due to ISM components not including dust, in percent, as described in §5.1 (Eq. 2b); (11) Same as Col. (10), but for the intrinsic LyC flux, i.e., the escape fraction of LyC including effects from all components of the ISM and reddening by dust.

the (blue) LyC aperture and a corresponding UVC measurement of $m_{\text{AB}}^{\text{UVC}} = 24.51$ mag (Table 2, Col. 5 and 12). The $\Delta m_{\text{AB}} = 5.58$ mag difference means that the LyC flux corresponds to $\langle F_{\text{UVC}}/F_{\text{LyC}} \rangle_{\text{obs}} = 10^{0.4 \cdot 5.58} \simeq 171 \pm 32$ when adding the relative errors in quadrature, or 162_{-17}^{+45} when taking these measurements as Gaussian distributions (see §5.2). Compared to the *intrinsic* LyC flux, the *intrinsic* stellar UVC flux of these galaxies increases by a factor $2.98_{-0.07}^{+0.08}$ (e.g., Siana et al. 2010). We determine this factor as expressed in Eq. 6 from the intrinsic SEDs and filter curves used for LyC and UVC observations, with their 1σ dispersion, as listed in Col. 4 of Table 3. These values suggest that the escape fraction for this particular case should be $\sim 2.98/162 \simeq 1.8\%$. Moreover, at $\langle z \rangle \simeq 2.68$, the *filter weighted* average IGM transmission of the redshifts in the stack is $0.247_{-0.085}^{+0.086}$, which was determined from the models of Inoue et al. (2014), and is listed in Col. 7 of Table 3. Hence, the $\mathcal{T}_{\text{IGM}}^{\text{LyC}}$ correction is a factor of $1/0.247 \simeq 4.05$, which requires the relative escape fraction of those LyC photons that make it out of the galaxies’ ISM to be on average $\sim 4.0 \times 1.8\% \simeq 7.4\% \pm 2.9$, or $6.0_{-1.0}^{+5.2}\%$ when assuming Gaussian photometric errors. This is the *directly calculated* $f_{\text{esc}}^{\text{rel}}$ value that is listed in Col. 8 of Table 3. These can be compared to Col. 9, which lists the more precise constraints on $f_{\text{esc}}^{\text{rel}}$ that incorporate the *wavelength-dependent* transmission models of LyC through the IGM (see §5.2).

The uncertainty in the direct $f_{\text{esc}}^{\text{rel}}$ calculations (in this particular case $\simeq 6.0_{-1.0}^{+5.2}\%$) fully propagate the SNR-induced errors in the measured LyC and UVC fluxes (Col. 5, 6 and 12, 13 of Table 2) and the $\pm 1\sigma$ errors on both the factor $\langle F_{\text{UVC}}/F_{\text{LyC}} \rangle_{\text{int}}$ and the $\langle \mathcal{T}_{\text{IGM}}^{\text{LyC}} \rangle$ values summarized in Table 3. We also include an estimate of the SED error from the 4–6 observed continuum data points that were used to fit each

SED at its fixed known redshift, i.e., the filters that sample the continuum emission of the galaxies longwards of Ly α up to WFC3/IR F125W, which are not affected by the IGM, even at high redshift. The main uncertainty in the SED fitting is therefore *not* the χ^2 values of fitted data, but the uncertainty in the applied internal extinction values A_V , which is unknown. We therefore do not include a dust correction *error* in our calculations. Calzetti et al. (2000) empirically derived the dust attenuation curves of nearby starburst galaxies and found a total to selective extinction value of $R_V = 4.05 \pm 0.8$. Our SEDs were reddened with various dust-screen A_V values by $F_{\nu}^{\text{obs}}(\lambda) = F_{\nu}^{\text{int}}(\lambda) 10^{-0.4 A_{\lambda}}$, where: $A_{\lambda} = k(\lambda) A_V / R_V$. The attenuation for wavelengths shorter than 630 Å and longer than 2200 Å were extrapolated from the interpolated slope of the endpoints of the attenuation curves. The applied reddening does not include the uncertainty in the R_V value. Estimating the reddening error in the flux of our sample galaxies would require a more extensive SED fitting analysis, which takes into account the equally probable A_V -values that fall within the measurement errors of the observed continuum data points. We instead assume a single A_V -value and vary the SED flux based on their observed continuum errors. This error estimate is equivalent to applying a convolution to the intrinsic SED LyC flux with a Gaussian kernel (see §5.2), and so applying an additional dust correction error would only increase the size of this kernel. Since the uncertainties in our measurements are dominated by the variation of the IGM transmission of LyC at the various redshifts and sight-lines, we did not attempt to quantify this additional dust-correction error. Nevertheless, our calculated f_{esc} -values themselves (§5.2) are not significantly affected by the A_V -uncertainty, though their quoted $\pm 1\sigma$ ranges would increase somewhat. Further details on the adopted A_V distributions are given in




Figure 11. Radial surface brightness profiles of the non-ionizing UVC signal (solid curves) and the LyC signal (dashed curves) measured in the stacks (Fig. 8) for the galaxies *without* AGN sample. The curves are color-coded according to their mean redshift (filter): $\langle z \rangle = 2.38$ (F606W and F225W; violet), $\langle z \rangle = 2.68$ (F606W and F275W; blue), $\langle z \rangle = 3.49$ (F775W and F336W; green), and $\langle z \rangle = 5.02$ (F850LP and F435W; red). The *observed* PSF in F275W and F435W is indicated by dotted blue and red curves, which were normalized to the central SB of the corresponding LyC surface brightness profiles. The horizontal black dashed line indicates the 1σ sensitivity limit for the LyC profile in F225W. All four UVC surface brightness profiles are extended with respect to the corresponding PSF curves. The observed LyC stack SB-profiles are also extended and flatter than the UVC profiles, which is also predicted from our LyC scattering model (light blue dashed curve; see §5.5), where scattering of the escaping LyC photons off electrons and/or dust with a porous ISM spreads the LyC emission beyond the distribution of the stellar UVC light (light blue solid curve).

Appendix D.1.

5.2. Estimating the LyC Escape Fraction: Monte Carlo Analysis

Since we cannot measure the amount of intrinsic LyC radiation produced by stars within galaxies directly, we must use the best available stellar population synthesis and IGM absorption models to estimate the fraction of LyC that escapes from galaxies at high redshift in a statistical way. With the best BC03 models fit to our 4–6 band photometric data at the known redshifts and the IGM transmission models of Inoue et al. (2014), we simulated the most likely f_{esc} values for the galaxies in our four redshift bins with a Monte Carlo analysis, treating the various measured and modeled fluxes and IGM transmission as discrete random variables (RVs), which incorporates a range of possible values for these measurements. This method also allows us to model the most likely apparent intrinsic flux from individual galaxies in a stack, which can vary substantially around the mean for galaxies at different redshifts observed through different IGM sight-lines (Inoue & Iwata 2008). We model all observed stacked fluxes from galaxies as Gaussian RVs, since the dominant sources of noise in the measurement (i.e., sky-background, shot- and read-noise) is approximately Gaussian.

For each galaxy, we take our best fit BC03 SED to obtain our dust-free (intrinsic) model LyC flux measurement

($F_{\nu, \text{LyC}, i}^{\text{int}}$), as denoted in Eq. 2a, before correcting for IGM absorption. We then approximate the error of this *model* LyC flux by independently varying the *observed* continuum data points the SEDs were fit to within their error bars, and re-fitting the SED by least squares, i.e., $F'_{\nu}(\nu) = \alpha F_{\nu}(\nu)$, where $\alpha = \mathbf{F}_{\nu}^r \cdot \mathbf{F}_{\nu}^m / \|\mathbf{F}_{\nu}^m\|^2$ and $\mathbf{F}_{\nu, i}^r$ and $\mathbf{F}_{\nu, i}^m$ are the randomized *observed* flux measurements and continuum band *model* fluxes, respectively. This process is equivalent to convolving the *model* LyC flux with a Gaussian kernel of width equal to the quadratic sum of the *observed* relative continuum errors. These continuum band *model* fluxes were calculated by convolving the apparent dust-attenuated SED with the respective filter transmission curve, similar to Eq. 4.

Using models of the IGM transmission with updated absorber statistics from Inoue et al. (2014), which simulates the transmission of photons through the IGM from $600\text{\AA} < \lambda_{\text{rest}} < 1300\text{\AA}$ for 10^4 lines-of-sight, we apply the IGM attenuation to our *model* LyC flux by convolving the SED with the wavelength dependent IGM transmission coefficient curve at the redshift of the galaxy for all 10^4 simulated lines-of-sight. We then convolve the IGM attenuated model flux with the model error that we calculated to obtain our final model LyC flux (i.e., $F_{\nu, \text{LyC}, i}^{\text{int}}$). We then stacked all of the IGM attenuated *model* LyC fluxes of all galaxies in their respective redshift bins to obtain our stacked *model* LyC flux as denoted in Eq. 2a. The stacked *model* LyC flux was then used to calculate $f_{\text{esc}}^{\text{abs}}$, $f_{\text{esc}}^{\text{rel}}$, and the dust attenuated $f_{\text{esc}}^{\text{abs}}$, as shown in Eqs. 2b–3a. Since we performed a non-correlated sum of the model LyC flux RVs to estimate this intrinsic stacked LyC flux, we run this f_{esc} calculation for a total of 10^3 trials, which we combine in order to generate a statistically significant sample of possible f_{esc} values. The probability mass function (PMF) of f_{esc} was then calculated by optimally binning these f_{esc} samples according to the Freedman-Diaconis rule (see §2.3), and normalizing by N_{samples} . The full f_{esc} PMFs are shown in Fig. 12 for galaxies without AGN. The statistics of the sample array, i.e., the ML values, averages, and $\pm 1\sigma$ error ranges were computed and are shown in Fig. 12 and Cols. (9)–(11) of Table 3.

Since each element in our sample represents a simulated possible value of f_{esc} , we take the mode, or the maximum-likelihood (ML) value of our PMF as the f_{esc} -value of our galaxies at their average redshift, as these values correspond to the escape fraction of the total flux from all galaxies in each stack. The upper error bars were computed as the $+1\sigma$ error (84%) of the full dataset, and the lower errors were calculated to be at the same 1σ likelihood value below the ML values. The averages and medians were computed from the full sample, and the lower bound on the averages was computed at -1σ (16%). The bin size of the $\langle z \rangle \simeq 5.02$ f_{esc} PMF was reduced to smaller than the Freedman-Diaconis prescription to obtain sub-percent resolution in the distribution. The various, less significant peaks in the $\langle z \rangle \simeq 2.38$ and $\langle z \rangle \simeq 2.68$ $f_{\text{esc}}^{\text{abs}}$ PMFs represent the apparent fluxes of fainter stellar populations that non-coherently contributed flux to the ML-model LyC flux. These *model* LyC fluxes also represent the lines-of-sight, where escaping LyC flux was transmitted through the IGM before being absorbed by Lyman Limit Sys-




Figure 12. [Left]: Probability mass functions (PMFs) of the *absolute* f_{esc} values from the MC simulations described in §5. [Right]: Same as left but with Calzetti et al. (2000) dust attenuation applied to the model LyC flux. This analysis was performed 10^3 times using the measured and modeled intrinsic stacked *apparent* LyC flux and their $\pm 1\sigma$ ranges. We apply the IGM attenuation models of Inoue et al. 2014 to our modeled LyC fluxes. These $f_{\text{esc}}^{\text{abs}}$ values were optimally binned according to the Freedman-Diaconis rule (see §4.1). Filled triangles and circles indicate the resulting ML and average f_{esc} values in each probability distribution function, respectively, while the horizontal tick marks indicate the $\pm 1\sigma$ range around the mode.

tems and Damped Lyman- α systems within $\Delta z \simeq 0.5$. The opaque lines-of-sight, where the IGM transmission peaks near $\mathcal{T}_{\text{IGM}}^{\text{LyC}} \simeq 0.01$, represent ~ 30 – 40% of our potential model LyC flux values. These lines-of-sight result in higher f_{esc} , as the model LyC would have been attenuated by more absorbers. However, ~ 40 – 50% of our lines-of-sight have average IGM transmission values $\mathcal{T}_{\text{IGM}}^{\text{LyC}} \gtrsim 0.4$ (where the transmission distribution is at a local minimum), and corresponds to the peak of the f_{esc} PMFs, where the model LyC flux encountered fewer absorbers. These lines-of-sight have a local maximum transmission near $\mathcal{T}_{\text{IGM}}^{\text{LyC}} \simeq 0.7$, and about $\sim 0.3\%$ can be as high as $\mathcal{T}_{\text{IGM}}^{\text{LyC}} \simeq 0.85$.

5.3. Implications of the f_{esc} MC Results

We list the results of our f_{esc} MC simulations in Table 3. The average absolute escape fraction, $\langle f_{\text{esc}}^{\text{abs}} \rangle$, from galaxies at various redshifts can be used to determine what fraction of LyC produced by the stellar photospheres in those galaxies escapes, i.e., is not absorbed by interstellar neutral H I, dust, etc., at their average redshift. However, variations in IGM transmission can cause these values to become highly uncertain when stacking LyC emission from galaxies at high and low redshifts. Thus, in order to ascertain any meaningful evolution in $f_{\text{esc}}^{\text{abs}}$, we must stack galaxies at similar redshifts and compare their $f_{\text{esc}}^{\text{abs}}$ values from sample to sample. Then, any trends in the independent subsamples can be used to constrain correlations of f_{esc} with galaxy properties or evolution with redshift. Modeling these properties can also be used to determine their impact on f_{esc} , and to see if trends in these properties with redshift can affect the apparent evolution of f_{esc} with cosmic time. For this reason, we explored the $f_{\text{esc}}^{\text{abs}}$ of galaxies both with and without dust attenuation applied, in order to understand the impact of dust on the $f_{\text{esc}}^{\text{abs}}$ values of these galaxies.

As seen in Table 3, the results of our f_{esc} analysis can vary by ~ 2 orders of magnitude when separating dust from the apparent SED of the galaxies (e.g., $\langle z \rangle = 2.380$, $\langle z \rangle = 2.682$). We also note that f_{esc} is not significantly affected when less dust is present in the majority of spectroscopically selected galaxies in a stack (as seems to be the case at $\langle z \rangle = 3.47$). However, the f_{esc} values when dust attention is applied only decreases by a few tens of percent, which indicates the percentage of LyC photons that is absorbed by components of the ISM that are *not* dust. Comparing the variations of f_{esc} from sample to sample, we see that dust is the dominant factor in reducing f_{esc} in galaxies at any redshift. This correlation of high dust extinction and low f_{esc} is consistent with the results of several observational and analytical studies that investigate the impact of various galactic parameters on f_{esc} (e.g., Mathis 1971; Leitherer et al. 1995; Inoue 2001; Bergvall et al. 2013).

The galaxies selected in our $\langle z \rangle \simeq 2.38$ and $\langle z \rangle \simeq 2.68$ stacks have, on average, younger stellar populations and more dust than the $\langle z \rangle \simeq 3.47$ and $\langle z \rangle \simeq 5.02$ stacks. The f_{esc} values for galaxies *selected* at $\langle z \rangle \simeq 3.47$ are indicative of older stellar populations (of ~ 1 Gyr), but is not significantly affected by the lower amount of dust observed in these galaxies. The $\langle z \rangle \simeq 2.38$ and $\langle z \rangle \simeq 2.68$ stacks sample galaxies that are undergoing a period of more active star-formation compared to the two higher redshift samples, which may have led to the accumulation of more dust in these galaxies, but also a brighter *intrinsic* LyC flux. The $\langle z \rangle \simeq 5.02$ stack is in general consistent with a younger stellar population and lower dust extinction. Thus, these $f_{\text{esc}}^{\text{abs}}$ values also imply that the ISM can absorb a larger fraction of LyC flux from older stellar populations than from younger ones when comparing $f_{\text{esc}}^{\text{abs}}$ from older and younger stellar populations. Although young stellar populations can produce more intrinsic LyC than older ones, which then has a higher probability of escaping the ISM,

dust can significantly reduce the efficiency of LyC escape by orders of magnitude more than interstellar hydrogen, as indicated by the decrease in $f_{\text{esc}}^{\text{abs}}$ when including effects from dust in Table 3.

5.4. The Observed Radial Surface Brightness-Profiles in UVC and LyC

The radial profiles of our LyC and UVC stacks from Fig. 8 are shown in Fig. 11. We construct all observed radial SB-profiles using the `ellipse` task (Busko 1996) in IRAF⁴. This package fits isophotal ellipses to an object’s flux distribution, given a set of initial parameters for the object for specified increments in radius. We start the initial elliptical isophotes at a semi-major axis radius of 5 pixels, centered on the centroids determined by SExtractor, after which the radii are linearly decreased/increased by 0.3 pixel. While the centers of the elliptical annuli are held fixed, the ellipticity and position angle are left as free parameters. The `ellipse` task also interpolates the fitted surface brightness profile to within the size of the central pixel, but we only retain a single data point, nominally at $r = 0''.045$, to represent our central SB and help normalize the corresponding PSFs.

The stacked UVC profiles are shown as solid curves, and those for LyC are dashed. We smoothed the LyC stacks by a $\sigma=1$ pixel Gaussian kernel before fitting. This smoothing did not affect the radial profiles substantially, as seen in Fig. 10, but reduced the noise in the stack so that the isophotes could be determined more accurately. The *observed* PSFs in the WFC3/UVIS F275W and ACS/WFC F435W mosaics are indicated by dotted curves, normalized to the central surface brightness of the corresponding LyC SB-profile. The PSFs in F225W and F336W are very similar to the F275W PSF, so we do not plot these (see Table 1 and Fig. 7b of W11). Note that these PSFs measured in the $0''.09$ mosaics are undersampled. The 1σ SB sensitivity limit for the LyC profile in the F225W stack is indicated by a horizontal dashed line at $\mu_{\text{AB}} \simeq 32.0$ mag arcsec⁻². For the sensitivity limits of the samples in the other filters, we refer the reader to Col. 9 of Table A1 and to the discussion of systematics in the stacked data in Appendices A–C. These SB sensitivity limits are consistent with the sky-subtraction errors discussed in §2.2. The sky-background in F435W is considerably brighter than for the three UV filters, since the Sun, and hence the Zodiacal sky-background, are significantly brighter longwards of the 4000Å break.

All four UVC SB-profiles are clearly extended with respect to their corresponding filter PSFs, as expected for stacked galaxy radial light-profiles at $z \simeq 3\text{--}6$ (e.g., Hathi et al. 2008). The much deeper HUDF UVC-stacks by Hathi et al. (2008) suggested a possible “break” (or slight change in slope) near $r \gtrsim 0''.3\text{--}0''.4$, from exponential in the inner parts to a somewhat less steep profile in the outskirts. Our stacked UVC light-profiles do not clearly show a change in slope at $r \gtrsim 0''.3\text{--}0''.4$, since our (77–180 orbit) UVC stacks are not nearly as deep as their $\sim 1680\text{--}4300$ orbit stacks, and because of our much more stringent method of masking neighbors.

⁴ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

The four LyC SB-profiles are also clearly extended with respect to their *observed* PSFs, and remain extended to $r \simeq 0''.7$ in F275W, F336W, and F435W, and to at least $r \sim 0''.4$ in F225W, beyond which errors in the sky-subtraction start to become substantial. The very faint, flat, non-centrally concentrated appearance of the combined LyC signal makes the extraction of its SB-profile uncertain at larger radii. The relatively flat LyC SB-profiles may indicate a more complicated LyC escape scenario, in which the light distribution of the LyC flux of a stack of galaxies is largely dependent on the porosity of the ISM in those galaxies, and/or the scattering processes that the LyC photons undergo before escape.

After integrating these SB-profiles as elliptical frustums between each isophote, we find reasonable agreement with our SExtractor photometry (see Table 2), although the flux represented by the radial SB profiles is consistently fainter by $\sim 0.3 \pm 0.2$ mag. This discrepancy is expected, given that our SB-profiles do not extend out to the aperture sizes used in the photometry of the stacks, and therefore miss some real LyC flux that might be present at larger radii and at very faint SB-levels.

5.5. Modeling the UVC and LyC Radial Surface Brightness Profiles

For the highest SNR measurements in the LyC stacks (i.e., F275W/F336W), the radial SB-profile of escaping LyC flux appears to be flatter than the corresponding *non-ionizing* UVC profile (the dashed and solid colored curves in Fig. 11, respectively). This appears to be the case somewhat less for the lower SNR F225W light-profile (purple dashed curve). The F435W stack has lower dynamic range in its SB-profile, as the sky brightness is higher in this filter compared to the three WFC3 UVIS filters.

A LyC SB-profile that is measurably flatter than the corresponding UVC SB profile could arise naturally in a porous ISM, in which the covering factor of neutral gas decreases with increasing galacto-centric distance. To illustrate this quantitatively, we consider the transfer of UVC and LyC continuum photons through simplified models of galaxies with a multiphase ISM.

To calculate this, we assume that the UVC sources are spatially extended and characterized by a volume emissivity $\epsilon_{\text{UVC}}(r)$. We assume an exponential distribution with galactic radius: $\epsilon_{\text{UVC}}(r) = \epsilon_{\text{UVC},0} \exp(-r/r_0)$. The normalization constant $\epsilon_{\text{UVC},0}$ and scale length of r_0 are obtained by matching the observed SB-profiles in Fig. 11. We further assume that LyC emission traces the UVC emission. We attribute differences in observed SB-profiles to the fact that neutral clumps of gas are opaque to LyC radiation, but not to UVC.

We also assume a (spherical) distribution of neutral gas clumps, which is described completely by its covering factor, $f_{\text{cov}}(r) \equiv n_c(r)A_c(r)$. Here, n_c and $A_c(r)$ denote the number of clumps and area of a clump at r , respectively. The covering factor f_{cov} then denotes the probability that a sight-line intersects a clump per unit length. For example, for clumps of fixed size that are outflowing at an *assumed constant* velocity v , we have a number density dependence as $f_{\text{cov}} \propto n_c \propto r^{-2}$ (we refer the reader to Dijkstra & Kramer (2012) for a more

detailed description of this covering factor).

The precise radial dependence of f_{cov} is not known. However, when f_{cov} decreases with r we generally expect increased LyC escape fractions at larger galacto-centric distances. We consider two-parameter models for $f_{cov} = Ar^{-x}$, and fit for A and x . Both A and x can also depend on radius. Hence, f_{cov} generally is some unknown power-law of r (i.e., r^{-x}), where x typically ranges between 0 and 3. This calculation shows that when sight-lines with low impact parameter see the largest f_{cov} , we see a reduced escape fraction in these directions. For example, at $\langle z \rangle = 2.68$ we find as best fit $(A, x) \simeq (0.11, 1.14)$.

An interesting possibility is that the neutral gas clouds can theoretically *scatter* LyC photons: LyC photons penetrate the neutral clumps over an average distance that corresponds to $\tau \simeq 1$. Direct recombination to the ground state produces LyC photons that can escape from the neutral cloud, as the optical depth to the edge of cloud is $\tau \simeq 1$. This “scattering” (absorption and re-emission of LyC photons occurs on the recombination time scale inside the cloud) of LyC photons would further flatten the predicted surface brightness profile, and can easily be included in the models of [Dijkstra & Kramer \(2012\)](#). The possible effects of LyC scattering can be expanded to also include scattering off free electrons and dust grains (which also differs between LyC and UVC).

While the dot-dashed curve in Fig. 11 is only a single example (matching our F275W LyC observations at $\langle z \rangle \simeq 2.68$) of these model LyC SB-profiles, model predictions with similar parameter values fit the SB profiles in the other redshift bins. With these models, we can integrate out to larger impact parameters and get a constraint on the total escape fraction that accounts for the difference in SB-profiles. This procedure gives a slightly larger value for f_{esc} than those given in Table 3 (This is, of course, almost identical to the constraints one would obtain simply by extrapolating the surface brightness profiles to larger impact parameters, and integrating over them.)

6. DISCUSSION OF RESULTS

6.1. Summary of Available Data on f_{esc} vs. Redshift

The relatively low values for f_{esc} that we find are valid for the luminosity range $M_{AB}^{UV} \simeq -20.8^{+0.9}_{-0.7}$ mag present in the sample selected to have reliable spectroscopic redshifts (see Fig. 5), with the two lowest redshift subsamples being dominated by relatively younger stellar populations with active star-formation and significant dust extinction, the $\langle z \rangle = 3.47$ subsample being comprised of mostly lower extinction galaxies with somewhat older stellar populations, and the $\langle z \rangle = 5.02$ sample including galaxies with younger stellar populations with lower dust extinction. For faint galaxies to have finished reionization by $z \simeq 6-7$, their f_{esc} values need to be $\gtrsim 20\%$ ([Ouchi et al. 2009](#); [Wilkins et al. 2011](#); [Kuhlen & Faucher-Giguère 2012](#)) beyond $z \simeq 6$ and their luminosities need to reach as faint as $M_{UV} \simeq -15$ to -13 mag (e.g., [Robertson et al. 2013](#)). Hence, if faint (dwarf) galaxies contributed significantly to reionization at $z \lesssim 6-7$, one should consider how much their f_{esc} fraction might have increased *both* towards higher redshifts due to the expected lower metallicities




Figure 13. The *absolute* LyC escape fraction for various galaxy samples as a function of redshift. Plotted are our ML average f_{esc}^{abs} values for our galaxies *without* AGN that have HQ spectra (orange filled circles) and their $\pm 1\sigma$ ranges taken from the probability mass functions of Fig. 12, generated from our MC simulations described in §5. The blue points indicate available published data as referenced in §1, some of which were converted from quoted f_{esc}^{rel} values using extinction values from the literature source (see §5.1, Eq. 3b). All vertical error bars are the $\pm 1\sigma$ uncertainty on the f_{esc} values. Some errors were converted from the quoted $2-3\sigma$ uncertainties. Upper and lower limits are shown as blue downward and upward triangles, respectively. Dark blue points indicate faint, nearby, star-bursting dwarf galaxies ($M_V \simeq -17.5$ mag). Although the blue points represent galaxy samples with different properties from our samples, and the quoted errors were derived from uncertainties with different error assessment, the combined data suggests a trend of f_{esc}^{abs} with redshift, which may not be a simple power law in $(1+z)$. For reference, we plot the best fit power law (purple dashed line): $f_{esc}^{abs} \simeq (0.56 \pm 0.21) \cdot (1+z)^{1.51 \pm 0.68} \%$ (68% confidence interval). We also plot a best fit sigmoidal curve in $(1+z)$ (green solid line), which had smaller residuals than the power law. This $\tanh(1+z)$ relation is fit to capture the observed sudden decrease in f_{esc}^{abs} from $f_{esc}^{abs} \simeq 20\%$ at $z \gtrsim 4$ to $f_{esc}^{abs} \sim 0.3\%$ at $z \lesssim 2$, with a midpoint at $z \simeq 3.1$ and $f_{esc}^{abs} \sim 11\%$. This relation implies that massive galaxies may have had high enough LyC f_{esc} values to complete hydrogen reionization by $z \sim 6$.

ties and lower dust extinction, *and* at fainter luminosities due to the larger impact that SN-driven outflows have on lower-mass dwarf galaxies (e.g., [Ricotti & Shull 2000](#); [Razoumov & SommerLarsen 2007](#); [Wise & Cen 2009](#); [Fernandez & Shull 2011](#)). Given that our spectroscopic selection samples luminous galaxies in all four redshift bins, our LyC detections can only constrain the first possibility, which we discuss here.

In Fig. 13, we plot our ML *average* f_{esc}^{abs} values generated from the MC simulation listed in Table 3 for galaxies without AGN activity (orange filled circles) along with the similarly derived f_{esc}^{abs} data available in published work summarized in §1 as blue points, with upper limits indicated as blue triangles. Light blue points indicate galaxies with restframe 1500Å luminosities close to those sampled in Fig. 5a–5c (i.e., $\langle M_{AB} \rangle \simeq -20.8 \pm 0.9$ mag). Dark blue points indicate faint, nearby, star-bursting dwarf galaxies of lower average luminosity ($\langle M_{AB} \rangle \simeq -17.5 \pm 0.3$ mag). The dependence of the f_{esc} -values of galaxies on luminosity is not well determined, but no clear dependence on luminosity is obvious in Fig. 12.

We will therefore address the redshift dependence of f_{esc} here for the luminosities sampled in Fig. 5.

We first converted the published $f_{\text{esc}}^{\text{rel}}$ values to $f_{\text{esc}}^{\text{abs}}$ when necessary using the quoted extinction values from the literature source. We note that these f_{esc} values were derived from different observational analyses, including both space and ground based spectra and imaging, with different object selection, reduction techniques, error assessment, and application of IGM attenuation models. Some of the published errors may not account for the same uncertainties that we address in §5.2. When necessary, we converted the quoted published uncertainties to 1σ error bars, so they are comparable to our results in Table 3.

Although the f_{esc} values plotted in Fig. 13 were derived with different methods, the current data appears to suggest a correlation of f_{esc} with redshift. However, this relation may not be a simple power-law in $(1+z)$. Several authors (Inoue et al. 2006; Razoumov & Sommer-Larsen 2010; Finlator et al. 2012; Kuhlen & Faucher-Giguère 2012; Becker & Bolton 2013; Dijkstra et al. 2014) have suggested that redshift-averaged f_{esc} values for galaxies may increase significantly with redshift, possibly as steeply as $\propto (1+z)^3 - (1+z)^5$. For reference, we fit a power law in $(1+z)$ to the available $f_{\text{esc}}^{\text{abs}}$ data in percent (Fig. 13), and plot the best fit (dashed purple line) as:

$$\langle f_{\text{esc}}^{\text{abs}} \rangle \simeq (0.56 \pm 0.21) \cdot (1+z)^{1.51 \pm 0.68}. \quad (7)$$

This only holds only for $z \lesssim 7$, beyond which the implied escape fraction would approach 100% for the upper bound, but decrease monotonically at lower redshift (e.g., Razoumov & Sommer-Larsen 2010). When applying the Anderson-Darling tests (Anderson & Darling 1952) to the $f_{\text{esc}}^{\text{abs}}$ data in Fig. 13, we find that the flatter power-law slope $\kappa \simeq 1.5$ of Eq. 7 is a better fit than published $f_{\text{esc}}(z)$ relations, which have steeper power-law slopes of $\kappa=3-5$. Since more than 1/3 of the data points falls outside the 68% confidence interval of this power-law fit, it is possible that the true $\langle f_{\text{esc}}^{\text{abs}} \rangle(z)$ relation would lie well below the lower bound of this power law. The center of the power law regression reaches $f_{\text{esc}}^{\text{abs}} \lesssim 10\%$ at $z \gtrsim 6-7$, which may *not* be sufficient for faint galaxies to finish reionization. If the $(1+z)^\kappa$ exponent values were as steep as $\kappa \simeq 2.0$, this regression would provide f_{esc} values at $z \gtrsim 6$ in excess of $\sim 30\%$, as required for hydrogen reionization to have completed by $z \sim 6$ (Robertson et al. 2013). However, none of the simple $(1+z)^\kappa$ power laws for f_{esc} are consistent with $\sim 2/3$ of the data points plotted in Fig. 13 to within their stated 1σ standard errors.

6.2. A Redshift Dependence Faster than $(1+z)^\kappa$?

Since more than half of the 49 independent data points in Fig. 13 deviates from the plotted best fit power law (violet dashed line) by more than 1σ , *no* single $(1+z)^\kappa$ regression seems to fit all the $f_{\text{esc}}^{\text{abs}}$ data for galaxies without AGN. We therefore suggest the possibility that a *more sudden decrease* of $f_{\text{esc}}^{\text{abs}}$ with redshift may instead have to be considered, as illustrated by the following “smooth-step”, or sigmoidal (“S”-

shaped) tanh-function in $(1+z)$:

$$\langle f_{\text{esc}}^{\text{abs}} \rangle \simeq \frac{\Delta}{2} \cdot \tanh([(1+z) - (1+z_0)]/\delta) + f_0, \quad (8)$$

where $\Delta \simeq 20.1 \pm 4.6$, $z_0 \simeq 3.13 \pm 0.04$, $\delta \simeq 0.22 \pm 0.20$, and $f_0 \simeq 10.3 \pm 2.3$ in percent (68% confidence intervals). This best fit tanh function is indicated by the solid green curve in Fig. 13, which has smaller residuals compared to the power-law fits (§6.1). Each of the four parameters in Eq. 8 can be interpreted as follows: Δ indicates the range of the $f_{\text{esc}}^{\text{abs}}$ values between their minima and maxima, z_0 indicates the pivot redshift of the tanh-function where the $f_{\text{esc}}^{\text{abs}}$ values decline most steeply with epoch, $\sim 5 \times \delta$ indicates over how wide a range in redshift the $f_{\text{esc}}^{\text{abs}}$ values drop from their maximum value at $z \gtrsim 4$ of $\sim 20\%$ to the lower values at $z \lesssim 2$ of $\sim 0.3\%$, and f_0 indicates the $f_{\text{esc}}^{\text{abs}}$ value at z_0 . Hence, Eq. 8 implies that massive galaxies may have had a sufficient f_{esc} of LyC to complete hydrogen reionization by $z \sim 6$. The combined data in Fig. 13 suggests, however, that $f_{\text{esc}}^{\text{abs}}$ may have declined by a factor of nearly ~ 100 from $\gtrsim 20\%$ at $z \gtrsim 3$ to $\sim 0.3\%$ at $z \lesssim 3$. These low f_{esc} -values at $z \lesssim 3$ are observed in some cosmological radiative transfer models as well, which also require a “steep-rise” in f_{esc} at $z \gtrsim 3$ for massive galaxies to reionize the Universe (e.g., Khaire et al. 2015), and have also been observed in studies of the Ly α escape fraction over redshift (e.g., Blanc et al. 2011). We note that we fit this “smooth-step” function solely to capture and quantify the observed steep decline in $f_{\text{esc}}^{\text{abs}}$, and that the true functional form of $f_{\text{esc}}^{\text{abs}}(z)$ is not yet uniquely constrained from the available data, especially at high redshift.

The width of the tanh function in $(1+z)$ indicates that the sudden decrease in $f_{\text{esc}}^{\text{abs}}$ may have occurred within the epoch of $z \sim 2.5-3.5$, or within about ± 0.8 Gyr of the observed peak in the cosmic star-formation history (SFH) (Madau et al. 1996; Faucher-Giguère et al. 2008; Cucciati et al. 2012; Burgarella et al. 2013). This period may indicate the epoch where the universe transitions from infall/merger driven star-forming galaxies at $3 \lesssim z \lesssim 6$ to a more passively evolving universe dominated by giant galaxies at $z \lesssim 1-2$ (Driver et al. 1998). This transition may have resulted in dust and gas rapidly accumulating in the disks and central bulges of forming galaxies, with a SN-rate that has progressively less impact on clearing gas and dust from the galaxies that are steadily growing in mass with cosmic time. It is therefore possible that this process may have caused $f_{\text{esc}}^{\text{abs}}$ to rapidly drop over a relatively narrow interval of cosmic time in luminous galaxies, as massive LyC-producing stars formed during the period of high SFR become either SNe II or AGB stars, which then enrich the ISM with dust within ~ 1 Gyr (Mathis 1990). The accumulated dust in these galaxies could have then caused $f_{\text{esc}}^{\text{abs}}$ to decrease substantially, and would also cause the dust to increase in temperature, as it absorbs most of LyC produced by the stars or AGN in the galaxy. This would then lead to a decrease in the galaxy’s SFR, as feedback from heating inhibits the formation of new massive stars (e.g., Inoue 2001; Inoue et al. 2001). The decline in SFR would also lead to a decreasing SN rate (Botticella et al. 2012), further preventing the escape of LyC, as there would have been fewer clear

channels produced by SN for the LyC to escape. LyC produced by AGN can be absorbed by gas and dust in the disk of the galaxy itself, depending on viewing angle. When galaxies produce stronger AGN outflows, more of their LyC radiation may escape approximately *orthogonally* from the galactic disk (e.g. Reunanen et al. 2003; Windhorst et al. 1998), which contributes to maintaining the ionized state of the IGM, as AGN begin to dominate the ionizing background at $z \lesssim 3$.

6.3. The Role of Galaxies with Weak AGN in Reionization

Fig 7 shows the stacked LyC and UVC images of the known galaxies with AGN in our sample. The $z=2.291$ stack only includes one AGN with a LyC flux of $m_{\text{AB}} \simeq 30.0$ mag at $\text{SNR} \sim 1$ (UVC aperture), so we do not remove the neighbors in the sub-image, but simply ignore these in the LyC photometry of this AGN by excluding them in the measurement aperture. The $\langle z \rangle \simeq 2.70$, 3.35, and 4.80 samples contain 7, 3, and 2 stacked AGN with measured LyC fluxes of $m_{\text{AB}} \simeq 28.9$, 29.7, and 28.6 mag with $\text{SNR} \sim 8.8$, 3.6, and 4.5, respectively. These fluxes are typically ~ 1 magnitude brighter than their non-AGN counterparts, despite having fewer stacked galaxies. This is most likely due to LyC originating from the central accretion disk, made visible by strong AGN outflows when viewed under the right angle. AGN outflows can also increase the porosity of the ISM in its host galaxy (e.g., Silk 2005), thereby increasing f_{esc} of the LyC produced by stars, which further contributes to the total measured LyC flux from that galaxy.

The stacks in Fig 7 show a variety of LyC morphologies, though the UVC images exhibit more compact light profiles compared to the non-AGN stacks in all three cases (see the discussion in §5.5–5.4). The $\langle z \rangle \simeq 2.70$ stack is the most extended of the AGN both in LyC and UVC, which is most likely due to the increased sensitivity to fainter flux at low redshift, with a central bright point-source from radiation escaping along the observed line-of-sight. The radial dependence of the LyC SB-profile for this stack may be due to the viewing angle of the AGN relative to the direction of the escaping LyC radiation, or possibly due to the fact that the LyC undergoes a more complex escape process, where photons can be reflected off of relativistic electrons in the AGN corona and accretion disk, or by hot dust in the torus via Thomson and/or inverse Compton scattering (e.g., Haardt & Maraschi 1993). The $\langle z \rangle \simeq 3.35$ AGN LyC stack appears to be more point-like, indicating that these observed LyC photons may be escaping predominantly along the line-of-sight, which is supported by the presence of broad emission lines in their spectra, although the more extended LyC emission may not be visible simply due to the average luminosity of these AGN at higher redshift dimming by an additional $\sim 40\%$ from $\langle z \rangle \simeq 2.70$ to $\langle z \rangle \simeq 3.35$. At $\langle z \rangle \simeq 4.80$, the AGN flux appears more elongated both in LyC and UVC compared to the two lower-redshift AGN stacks, which is dominated by the contribution from an AGN at $z=4.823$. The X-ray counterpart for the $z=4.823$ AGN, which was used to identify this AGN (see Fiore et al. 2012), is also elongated in the same direction, which may indicate that the observed flux is the products of multiple inverse Compton scatterings of photons from longer

wavelengths into the FUV and X-ray, which then escapes in collimated cones approximately orthogonal to the accretion disk (e.g., Zheng et al. 1997; Telfer et al. 2002; Shull et al. 2012).

Fig. 5 shows that the *average* UVC luminosities of “Galaxies with weak AGN” in our sample is about the same, or somewhat fainter than that of galaxies without AGN. Their average luminosity in Fig. 5 $M_{\text{AB}} \simeq -20.4 \pm 0.9$ mag at $z \simeq 2.3\text{--}5$ does not indicate clearly QSO-dominated luminosities or SEDs. Table 2 shows that the LyC flux measured from the stacks of (weak) AGN at all redshifts is typically ~ 1 mag *brighter* than galaxies without AGN. Thus, precise modeling of the intrinsic LyC emission must include the contribution of flux emitted by, or reprocessed from, the AGN accretion disk. The SED of the AGN accretion disk may be more complicated than a simple blackbody curve, as the SED must account for the broad and narrow emission line regions, as well as energy lost to relativistic jets and photons scattered/absorbed by the corona and central torus and non-AGN dust, which is also viewing-angle dependent. Because we cannot fit both stellar+AGN SED models to the 4–6 band continuum data currently available for galaxies with weak AGN, we do not calculate escape fractions for these galaxies. Since the SEDs of these galaxies are likely dominated by stellar light at the non-ionizing wavelengths, their $f_{\text{esc}}(z)$ correlation may be similar to the function derived from the escape fractions of galaxies without AGN. From the compact appearance in some of our stacked images — and from the fact that they are on average ~ 1 mag brighter than galaxies without AGN — the LyC flux in galaxies with weak AGN may be dominated by light originating from their accretion disks.

Further data and modeling is needed to better constrain $f_{\text{esc}}^{\text{abs}}(z)$ for both galaxies and weak AGN to confirm these observed trends. The current data for AGN may be consistent with a more modest drop in $f_{\text{esc}}^{\text{abs}}(z)$ than for galaxies that may have occurred close to the peak in the epoch of AGN activity around $z \simeq 2.5$ (e.g., Fontanot et al. 2007; Croom et al. 2009; Ikeda et al. 2011, 2012). Since AGN activity can affect the SFRs, it is possible that when AGN-outflows started to ramp up after the peak in the cosmic star-formation history at $z \simeq 3$ (Springel et al. 2005; Hopkins et al. 2006), their outflows cleared enough paths in the host galaxy ISM to increase $f_{\text{esc}}^{\text{abs}}$ of a possibly AGN-induced top-heavy stellar population IMF.

Because galaxies far outnumber AGN, and despite being fainter in LyC by ~ 1 mag on average, their $f_{\text{esc}}^{\text{abs}}$ values suggest that galaxies may have produced sufficient LyC radiation to maintain reionization at $z \gtrsim 3$, while AGN likely dominated in the production of ionizing LyC flux at $z \lesssim 3$.

The current samples are still very small, and clearly need further confirmation through much larger samples, both through deep UV/optical imaging of wider HST fields and through spectroscopy on fields with high quality existing HST data. Further theoretical work is needed to outline exactly how quickly f_{esc} may have increased towards higher redshifts *and* at fainter luminosities, as well as at lower metallicities and lower extinction at higher redshifts, while producing enough escaping LyC photons from faint galaxies to finish

and maintain reionization at $z \lesssim 6\text{--}7$.

There is already a significant issue in accounting for reionization with the faint galaxy population observed via cluster lensing at $z \simeq 9$. At redshifts larger than 8, the Hubble Frontier Fields reveal a strong drop in rest-frame UV luminosity density (e.g., Ishigaki et al. 2015). This issue may be compounded by our relatively low measured values for $f_{\text{esc}}^{\text{abs}}$ of $\sim 0.3\%$ for $M_{\text{AB}} \simeq -20.8$ mag at $z \simeq 2.3\text{--}2.7$. Hence, it is also possible that one may need to consider an additional source of reionizing photons beyond $z \simeq 6\text{--}7$. This source might include feedback on both the IGM ionization and clumpiness via hard ionizing photons from high mass X-ray binaries (e.g., Mirabel et al. 2011). Other astrophysical sources such as Population III stars or mini-AGN seem strongly constrained via chemical evolution (Kulkarni et al. 2014) and the X-ray background (Dijkstra et al. 2004). It is possible that f_{esc} may evolve with redshift and/or with galaxy properties (e.g., mass, A_V , SFR, and/or age). If such dependencies did not occur, sources of reionization other than faint galaxies may need to be considered.

7. CONCLUSIONS

We studied LyC emission that may be escaping from galaxies using improved *HST* WFC3 and ACS mosaics of the ERS and GOODS-S fields in four filters, where LyC may be observed from galaxies at $z \simeq 2.3\text{--}5$. The data that we used in our analysis was drizzled with the much more accurate 2013 WFC3 geometric distortion correction tables, which resulted in the correction of significant astrometric offsets that remained in earlier ERS UVIS mosaics. The WFC3 ERS UV images were taken in 2009 September, when the CTE was still at a level where faint flux could still be measured without significant losses. We verified that any loss in CTE is not the primary limitation to our measurements (see Appendix C.2).

We extracted sub-images centered on galaxies with high quality spectroscopically measured redshifts from the ERS mosaics, and averaged the LyC flux of those galaxies. We paid careful attention to the removal of potentially nearby contaminating objects and low-level variations in the UV sky-background during this stacking process. We ensured that no significant amount of contaminating flux longwards of the Lyman-break ($\lambda > 912\text{\AA}$) was included in our stacks. We performed a series of critical tests to ensure our stacking procedure was not affected by various systematics in the mosaic images. All of these tests are described in Appendix A–C. The following are our main findings:

- (1) Our measurements of the average LyC flux that escapes from galaxies at $z \simeq 2.4\text{--}5$ is summarized in Table 2. We find that the LyC signal of faint galaxies at $\langle z \rangle \simeq 2.38, 2.68, 3.47$ and 5.02 is generally measured at the $\gtrsim 3\text{--}6\sigma$ level, in typical image stacks of 11–15 objects in the WFC3/UVIS F225W, F275W, F336W, and ACS/WFC F435W filters, respectively. Their measured signal corresponds to total LyC fluxes of $m_{\text{AB}} \simeq 29.5\text{--}30.7$ mag. The LyC flux of weak AGN is detected at levels $\sim 1 \pm 0.4$ mag brighter on average at $z \simeq 2.3\text{--}4.8$, but over $\sim 2\text{--}8\times$ fewer objects per stack.
- (2) The combined LyC emission averaged over the four filters suggests an overall LyC flux distribution that is non-

centrally concentrated, which may be explained by a radial dependence in the ISM porosity and/or scattering of the LyC photons. We find that the LyC flux from galaxies is more extended than its UVC counterpart in all four filters. This suggests a complex escape process that may be determined by the distribution and extent of neutral (dusty) gas clouds within a porous multiphase ISM.

- (3) From our best-fit BC03 SED models fit to *HST* continuum observations longwards of $\text{Ly}\alpha$, the observed LyC flux corresponds to an *average* absolute LyC escape fraction that increases from $f_{\text{esc}}^{\text{abs}} \sim 0.1\%$ at $\langle z \rangle \simeq 2.4\text{--}2.7$ to $\sim 18\%$ at $\langle z \rangle \simeq 3.5$. It is $\sim 4\%$ (with a very large error range) at $\langle z \rangle \simeq 5.0$, where we do see some LyC flux at the $\gtrsim 3\sigma$ level. The implied f_{esc} -value at $\langle z \rangle \simeq 5.0$ is very poorly constrained due to the considerable variation in the IGM attenuation over this redshift range and the sight-lines at $z \gtrsim 5$. We verify these MC derived f_{esc} values, which model the apparent intrinsic and dust-attenuated LyC fluxes with their $\pm 1\sigma$ errors through 10^4 random lines-of-sight, with “direct” calculations using our LyC–UVC photometry and average IGM transmission models of Inoue et al. (2014). While the error bars on the implied f_{esc} values in each of the four redshift bins remain large, within the error bars, the data suggest an increasing trend of f_{esc} with redshift at $z \gtrsim 3$.

- (4) The available published f_{esc} data for galaxies is bounded by $f_{\text{esc}}^{\text{abs}} \simeq (0.6 \pm 0.2) \cdot (1+z)^{1.5 \pm 0.7}$ (68% confidence interval). A better description seems to be a *more sudden increase* in $f_{\text{esc}}^{\text{abs}}$ with redshift that occurred around $z \sim 3.1$. Our best fits of $f_{\text{esc}}(z)$ to the available data are a simple non-linear sigmoidal (“S”-shaped) function in $(1+z)$, where each of the parameters has a simple physical meaning. For galaxies, the steepest drop in f_{esc} occurs at $z \simeq 3.1$, near the peak of the cosmic star-formation history within an interval of ± 0.8 Gyr from this peak in cosmic time. The pivot point for galaxies at $z \simeq 3.1$ occurs at $f_0 \simeq 11\%$, while the combined data suggest that $f_{\text{esc}}^{\text{abs}}$ may have dropped by nearly a factor of 100 from $\sim 20\%$ at $z \gtrsim 3$ to $\sim 0.3\%$ at $z \lesssim 3$ over a relatively short interval in cosmic time (~ 1.5 Gyr).

- (5) Because galaxies far outnumber weak AGN (by a factor of $\sim 2\text{--}8$ in our samples), and from their measured $f_{\text{esc}}^{\text{abs}}$ values, galaxies may have had a sufficient LyC f_{esc} to reionize the IGM by $z \gtrsim 6$. The SEDs of galaxies with weak AGN is likely dominated by *stellar light* in the non-ionizing continuum. However, their observed LyC flux outshines galaxies *without* AGN in our sample by $m_{\text{AB}} \sim 1$ mag. Hence, while galaxies without AGN likely began and maintained cosmic reionization at $z \gtrsim 3$, galaxies with (weak) AGN likely dominated the contribution to the cosmic ionizing background and maintain reionization at $z \lesssim 3$, although the role of massive galaxies without AGN may not have been negligible at $z \lesssim 3$.

The transition from galaxy-dominated reionization to weak AGN-dominated reionization appears to have occurred at $z \sim 2\text{--}3$, i.e., right around the peak in the cosmic SFR (Madau et al. 1996), which may indicate the epoch where the universe transitions from infall/merger driven SFGs at $3 \lesssim z \lesssim 6$ to a more passively evolving universe dominated by giant galaxies at $z \lesssim 1\text{--}2$. This may result in dust and gas rapidly accumulating in the disks and nuclei of forming galaxies, combined

with a SN-rate that has progressively less impact on clearing gas/dust in galaxies that are steadily growing in mass with cosmic time. The accumulating dust and decreasing SFR may have caused $f_{\text{esc}}^{\text{abs}}$ to *rapidly drop* over a relatively narrow interval of cosmic time (~ 1.5 Gyr), as the LyC flux heats the dust and inhibits the formation of new massive stars. When AGN-outflows began to increase after the peak in the cosmic star-formation history at $z \sim 3$, their outflows may have cleared enough paths in the ISM of host galaxies to enhance the fraction of escaping LyC radiation produced by massive stars and from the accretion disk, resulting in AGN beginning to dominate the ionizing background at $z \lesssim 3$.

• (6) Further data on LyC f_{esc} are essential for both galaxies and weak AGN to confirm both their trends in $f_{\text{esc}}^{\text{abs}}(z)$. The current samples are still very small, and clearly need further confirmation through much larger samples, both through deep imaging of wider HST fields in the UV and through deeper spectroscopy on fields with high quality existing HST data, e.g., with the *JWST* FGS/NIRISS grisms and with NIRSpec (Gardner et al. 2006). Further theoretical work is needed to outline exactly how quickly f_{esc} may have increased towards higher redshifts and at fainter luminosities, as well as at lower metallicities and lower dust extinction at higher redshifts, while producing enough escaping LyC photons from faint galaxies to complete and maintain reionization at $z \sim 6-6.5$.

ACKNOWLEDGMENTS

This paper is based on Early Release Science observations made by the WFC3 Scientific Oversight Committee. We are grateful to the Director of the Space Telescope Science Institute, Dr. Matt Mountain, for generously awarding Director's Discretionary time for this program. Finally, we are deeply indebted to the crew of STS-125 for refurbishing and repairing *HST*. We thank Drs. George Becker, Renyue Cen, Nimish Hathi, Anne Jaskot, Karen Olson, and Michael Rutkowski, and Mr. Jacob Vehonsky for their useful comments and suggestions for this work. Support for *HST* programs GO-11359 and AR-13877 was provided by NASA through grants from the STScI, which is operated by the Association of Universities for Research Inc., under NASA contract NAS 5-26555.

REFERENCES

- Ade, P. A. R., Aghanim, N., Armitage-Caplan, C., et al. 2014, *A&A*, 571, A16
- Aird, J., Coil, A. L., Georgakakis, A., et al. 2015, *MNRAS*, 451, 1892
- Anderson, J., & Bedin, L. R. 2010, *PASP*, 122, 1035
- Anderson, T. W., & Darling, D. A. 1952, *The Annals of Mathematical Statistics*, 23, 193
- Avila, R., Grogg, N., Anderson, J., et al. 2015, *ACS Instrument Handbook* (STScI, Baltimore), Version 14
- Baggett, S., & Anderson, J. 2012, *WFC3/UVIS Sky Backgrounds*, Tech. rep.
- Baggett, S., Brown, T., Boucarut, R., et al. 2006, in *Space Telescopes and Instrumentation I: Optical*, ed. J. C. Mather, H. A. MacEwen, & M. W. M. de Graauw, Vol. 6265, 626532–626532–11
- Balestra, I., Mainieri, V., Popesso, P., et al. 2010, *A&A*, 512, A12
- Becker, G. D., & Bolton, J. S. 2013, *MNRAS*, 436, 1023
- Becker, G. D., Bolton, J. S., Madau, P., et al. 2015, *MNRAS*, 447, 3402
- Bergvall, N., Leitert, E., Zackrisson, E., & Marquart, T. 2013, *A&A*, 554, A38
- Bergvall, N., Zackrisson, E., Andersson, B.-G., et al. 2006, *A&A*, 448, 513
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bielby, R., Hill, M. D., Shanks, T., et al. 2013, *MNRAS*, 430, 425
- Blanc, G. A., Adams, J. J., Gebhardt, K., et al. 2011, *ApJ*, 736, 31
- Bland-Hawthorn, J., & Maloney, P. R. 1999, *ApJ*, 510, L33
- Borthakur, S., Heckman, T. M., Leitherer, C., & Overzier, R. A. 2014, *Science*, 346, 216
- Botticella, M. T., Smartt, S. J., Kennicutt, R. C., et al. 2012, *A&A*, 537, A132
- Bourque, M., & Kozhurina-Platais, V. 2013, *Instrument Science Report WFC 2013-03* (STScI, Baltimore)
- Boutsia, K., Grazian, A., Giallongo, E., et al. 2011, *ApJ*, 736, 41
- Bouwens, R. J., Illingworth, G. D., Oesch, P. A., et al. 2012, *ApJL*, 752, L5
- Bridge, C. R., Teplitz, H. I., Siana, B., et al. 2010, *ApJ*, 720, 465
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Burgarella, D., Buat, V., Gruppioni, C., et al. 2013, *A&A*, 554, A70
- Busko, I. C. 1996, *Astronomical Data Analysis Software and Systems V*, 101
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, 533, 682
- Cen, R., & Kimm, T. 2015, *ApJ*, 801, L25
- Chen, H.-W., Prochaska, J. X., & Gnedin, N. Y. 2007, *ApJL*, 667, L125
- Choudhury, T. R., Puchwein, E., Haehnelt, M. G., & Bolton, J. S. 2015, *MNRAS*, 452, 261
- Clarke, C., & Oey, M. S. 2002, *MNRAS*, 337, 1299
- Cooke, J., Ryan-Weber, E. V., Garel, T., & Diaz, C. G. 2014, *MNRAS*, 441, 837
- Cowie, L. L., Barger, A. J., & Trouille, L. 2009, *ApJ*, 692, 1476
- Cristiani, S., Appenzeller, I., Arnouts, S., et al. 2000, *A&A*
- Croom, S. M., Richards, G. T., Shanks, T., et al. 2009, *MNRAS*, 399, 1755
- Cucciati, O., Tresse, L., Ilbert, O., et al. 2012, *A&A*, 539, A31
- de Barros, S., Vanzella, E., Amorin, R., et al. 2015, *ArXiv e-prints*, arXiv:1507.06648
- Deharveng, J.-M., Buat, V., Le Brun, V., et al. 2001, *A&A*, 375, 805
- Dickinson, M., Giavalisco, M., & GOODS Team. 2003, *The Mass of Galaxies at Low and High Redshift*, ed. R. Bender & A. Renzini, *ESO ASTROPHYSICS SYMPOSIA* (Berlin/Heidelberg: Springer-Verlag), doi:10.1007/b10204
- Dijkstra, M., Haiman, Z., & Loeb, A. 2004, *ApJ*, 613, 646
- Dijkstra, M., & Kramer, R. 2012, *MNRAS*, 424, 1672
- Dijkstra, M., Mesinger, A., & Wyithe, J. S. B. 2011, *MNRAS*, 414, 2139
- Dijkstra, M., Wyithe, S., Haiman, Z., Mesinger, A., & Pentericci, L. 2014, *MNRAS*, 440, 3309
- Dove, J. B., Shull, J. M., & Ferrara, A. 2000, *ApJ*, 531, 846
- Dressel, L., Balick, B., Baggett, S., et al. 2015, *Wide Field Camera 3 Instrument Handbook* (STScI, Baltimore), Version 7.
- Driver, S. P., Fernández-Soto, A., Couch, W. J., et al. 1998, *ApJL*, 496, L93
- Ebrero, J., Carrera, F. J., Page, M. J., et al. 2009, *A&A*, 493, 55
- Fan, X., Carilli, C., & Keating, B. 2006a, *ARA&A*, 44, 415
- Fan, X., Narayanan, V. K., Strauss, M. A., et al. 2002, *AJ*, 123, 1247
- Fan, X., Narayanan, V. K., Lupton, R. H., et al. 2001, *AJ*, 122, 2833
- Fan, X., Strauss, M. A., Richards, G. T., et al. 2006b, *AJ*, 131, 1203
- FaucherGiguère, C., Lidz, A., Hernquist, L., & Zaldarriaga, M. 2008, *ApJ*, 688, 85
- Ferguson, A. M. N., Wyse, R. F. G., Gallagher, J. S., I., & Hunter, D. A. 1996, *AJ*, 111, 2265
- Ferguson, H. C. 2001, in *Deep Fields*, ed. S. Cristiani, A. Renzini, & R. E. Williams, 54
- Fernández, E. R., & Shull, J. M. 2011, *ApJ*, 731, 20
- Fernández-Soto, A., Lanzetta, K. M., & Chen, H.-W. 2003, *MNRAS*, 342, 1215
- Finkelstein, S. L., Ryan, Jr., R. E., Papovich, C., et al. 2015, *ApJ*, 810, 71
- Finlator, K., Oh, S. P., Özel, F., & Davé, R. 2012, *MNRAS*, 427, 2464
- Fiore, F., Puccetti, S., Grazian, A., et al. 2011, *A&A*, 537, A16
- Fiore, F., Puccetti, S., Grazian, A., et al. 2012, *A&A*, 537, A16
- Fontanot, F., Cristiani, S., Monaco, P., et al. 2007, *A&A*, 461, 39

- Freedman, D., & Diaconis, P. 1981, *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 57, 453
- Fruchter, A. S., et al. 2010, 2010 Space Telescope Science Institute Calibration Workshop, 382
- Fujita, A., Martin, C. L., Mac Low, M., & Abel, T. 2003, *ApJ*, 599, 50
- Fynbo, J. P. U., Jakobsson, P., Prochaska, J. X., et al. 2009, *ApJS*, 185, 526
- Gardner, J. P., Mather, J. C., Clampin, M., et al. 2006, *Space Sci. Rev.*, 123, 485
- Giallongo, E., Cristiani, S., D’Odorico, S., & Fontana, A. 2002, *ApJ*, 568, L9
- Giallisco, M., Ferguson, H. C., Koekemoer, A. M., et al. 2004, *ApJ*, 600, L93
- Gil de Paz, A., Boissier, S., Madore, B. F., et al. 2007, *ApJS*, 173, 185
- Gnedin, N. Y. 2000, *ApJ*, 535, 530
- . 2008, *ApJ*, 673, L1
- Gonzaga, S., et al. 2012, *The DrizzlePac Handbook* (STScI, Baltimore)
- Grazian, A., Giallongo, E., Gerbasi, R., et al. 2016, *A&A*, 585, A48
- Grimes, J. P., Heckman, T., Strickland, D., et al. 2007, *ApJ*, 668, 891
- Grimes, J. P., Heckman, T., Aloisi, A., et al. 2009, *ApJS*, 181, 272
- Grogan, N. A., Kocevski, D. D., Faber, S. M., et al. 2011, *ApJS*, 197, 35
- Guaita, L., Pentericci, L., Grazian, A., et al. 2016, *ArXiv e-prints*, arXiv:1601.03057
- Guo, Y., Ferguson, H. C., Giallisco, M., et al. 2013, *ApJS*, 207, 24
- Haardt, F., & Madau, P. 1996, *ApJ*, 461, 20
- . 2012, *ApJ*, 746, 125
- Haardt, F., & Maraschi, L. 1993, *ApJ*, 413, 507
- Hanish, D. J., Oey, M. S., Rigby, J. R., de Mello, D. F., & Lee, J. C. 2010, *ApJ*, 725, 2029
- Hathi, N. P., Jansen, R. A., Windhorst, R. A., et al. 2008, *AJ*, 135, 156
- Hathi, N. P., Ryan, R. E., Cohen, S. H., et al. 2010, *ApJ*, 720, 1708
- Heckman, T. M., Sembach, K. R., Meurer, G. R., et al. 2001, *ApJ*, 558, 56
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2013, *ApJS*, 208, 19
- Hopkins, P. F., Somerville, R. S., Hernquist, L., et al. 2006, *ApJ*, 652, 864
- Hunt, M. P., Steidel, C. C., Adelberger, K. L., & Shapley, A. E. 2004, *ApJ*, 605, 625
- Hurwitz, M., Jelinsky, P., & Dixon, W. V. D. 1997, *ApJ*, 481, L31
- Ikeda, H., Nagao, T., Matsuoka, K., et al. 2011, *ApJ*, 728, L25
- . 2012, *ApJ*, 756, 160
- Inoue, A. K. 2001, *AJ*, 122, 1788
- Inoue, A. K., Hirashita, H., & Kamaya, H. 2001, *ApJ*, 555, 613
- Inoue, A. K., & Iwata, I. 2008, *MNRAS*, 387, 1681
- Inoue, A. K., Iwata, I., & Deharveng, J.-M. 2006, *MNRAS*, 371, L1
- Inoue, A. K., Iwata, I., Deharveng, J.-M., Buat, V., & Burgarella, D. 2005, *A&A*, 435, 471
- Inoue, A. K., Shimizu, I., Iwata, I., & Tanaka, M. 2014, *MNRAS*, 442, 1805
- Ishigaki, M., Kawamata, R., Ouchi, M., et al. 2015, *ApJ*, 799, 12
- Iwata, I., Inoue, A. K., Matsuda, Y., et al. 2009, *ApJ*, 692, 1287
- Izotov, Y. I., Orlitova, I., Schaerer, D., et al. 2016, *ArXiv e-prints*, arXiv:1601.03068
- Jensen, H., Laursen, P., Mellema, G., et al. 2012, *MNRAS*, 428, 1366
- Khair, V., Srianand, R., Choudhury, T. R., & Gaikwad, P. 2015, *ArXiv e-prints*, arXiv:1510.04700
- Koekemoer, A. M., Faber, S. M., Ferguson, H. C., et al. 2011, *ApJS*, 197, 36
- Koekemoer, A. M., Ellis, R. S., McLure, R. J., et al. 2013, *ApJS*, 209, 3
- Kozhurina-Platais, V. 2014, *Instrument Science Report WFC 2014-12* (STScI, Baltimore)
- Kozhurina-Platais, V., Cox, C., McLean, B., et al. 2009, *Instrument Science Report WFC3 2009-33* (STScI, Baltimore)
- Kozhurina-Platais, V., Hammer, D., Dencheva, N., & Hack, W. 2013, *Instrument Science Report WFC 2013-14* (STScI, Baltimore)
- Kuhlen, M., & Faucher-Giguère, C.-A. 2012, *MNRAS*, 423, 862
- Kulkarni, G., Hennawi, J. F., Rollinde, E., & Vangioni, E. 2014, *ApJ*, 787, 64
- Kurk, J., Cimatti, A., Daddi, E., et al. 2012, *A&A*, 549, A63
- Le Fèvre, O., Vettolani, G., Paltani, S., et al. 2004, *A&A*, 428, 1043
- Leitet, E., Bergvall, N., Hayes, M., Linné, S., & Zackrisson, E. 2013, *A&A*, 553, A106
- Leitet, E., Bergvall, N., Piskunov, N., & Andersson, B.-G. 2011, *A&A*, 532, A107
- Leitherer, C., Ferguson, H. C., Heckman, T. M., & Lowenthal, J. D. 1995, *ApJ*, 454, doi:10.1086/309760
- Leitherer, C., Vacca, W. D., Conti, P. S., et al. 1996, *ApJ*, 465, 717
- Lilly, S., Schade, D., Ellis, R., et al. 1998, *ApJ*, 500, 75
- Loeb, A., & Barkana, R. 2001, *ARA&A*, 39, 19
- Lusso, E., Worseck, G., Hennawi, J. F., et al. 2015, *MNRAS*, 449, 4204
- Mack, J., Sabbi, E., & Dahlen, T. 2013, *Instrument Science Report WFC3 2013-10* (STScI, Baltimore)
- Madau, P. 1999, in *The 9th astrophysics conference: After the dark ages, when galaxies were young (the Universe at $2 < z < 5$)*, Vol. 470 (ASCE), 299–311
- Madau, P., Ferguson, H. C., Dickinson, M. E., et al. 1996, *MNRAS*, 283, 1388
- Madau, P., & Haardt, F. 2015, *ApJL*, 813, L8
- Madau, P., Rees, M. J., Volonteri, M., Haardt, F., & Oh, S. P. 2004, *ApJ*, 604, 484
- Malkan, M., Webb, W., & Konopacky, Q. 2003, *ApJ*, 598, 878
- Mallén-Ornelas, G., Lilly, S. J., Crampton, D., & Schade, D. 1999, *ApJ*, 518, L83
- Massey, R., Schrabback, T., Cordes, O., et al. 2014, *MNRAS*, 439, 887
- Mathis, J. S. 1971, *ApJ*, 167, 261
- Mathis, J. S. 1990, *ARA&A*, 28, 37
- McGreer, I. D., Mesinger, A., & D’Odorico, V. 2014, *MNRAS*, 447, 499
- Mesinger, A., Aykutalp, A., Vanzella, E., et al. 2014, *MNRAS*, 446, 566
- Mesinger, A., & Haiman, Z. 2004, *ApJ*, 611, L69
- Miller, N. A., Bonzini, M., Fomalont, E. B., et al. 2013, *ApJS*, 205, 13
- Mirabel, I. F., Dijkstra, M., Laurent, P., Loeb, A., & Pritchard, J. R. 2011, *A&A*, 528, A149
- MiraldaEscudé, J., Haehnelt, M., & Rees, M. J. 2000, *ApJ*, 530, 1
- Moos, H. W., Cash, W. C., Cowie, L. L., et al. 2000, *ApJ*, 538, L1
- Mostardi, R. E., Shapley, A. E., Nestor, D. B., et al. 2013, *ApJ*, 779, 65
- Mostardi, R. E., Shapley, A. E., Steidel, C. C., et al. 2015, *ApJ*, 810, 107
- Nestor, D. B., Shapley, A. E., Kornei, K. A., Steidel, C. C., & Siana, B. 2013, *ApJ*, 765, 47
- Nestor, D. B., Shapley, A. E., Steidel, C. C., & Siana, B. 2011, *ApJ*, 736, 18
- Noeske, K., Baggett, S., Bushouse, H., et al. 2012, *WFC3 UVIS Charge Transfer Efficiency October 2009 to October 2011*, Tech. rep.
- Oey, M., & Kennicutt, R. C., J. 1997, *MNRAS*, 291
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Ouchi, M., Mobasher, B., Shimasaku, K., et al. 2009, *ApJ*, 706, 1136
- Planck Collaboration. 2015, eprint arXiv:1502.01589
- Popesso, P., Dickinson, M., Nonino, M., et al. 2009, *A&A*, 494, 443
- Putman, M. E., BlandHawthorn, J., Veilleux, S., et al. 2003, *ApJ*, 597, 948
- Rafelski, M., Teplitz, H. I., Gardner, J. P., et al. 2015, *AJ*, 150, 31
- Rangel, C., Nandra, K., Laird, E. S., & Orange, P. 2013, *MNRAS*, 428, 3089
- Razoumov, A. O., & Sommer-Larsen, J. 2010, *ApJ*, 710, 1239
- Razoumov, A. O., & SommerLarsen, J. 2007, *ApJ*, 668, 674
- Reddy, N. A., & Steidel, C. C. 2009, *ApJ*, 692, 778
- Reunanen, J., Kotilainen, J. K., & Prieto, M. A. 2003, *MNRAS*, 343, 192
- Ricotti, M. 2002, *MNRAS*, 336, L33
- Ricotti, M., & Shull, J. M. 2000, *ApJ*, 542, 548
- Robertson, B. E., Furlanetto, S. R., Schneider, E., et al. 2013, *ApJ*, 768, 71
- Rutkowski, M. J., Scarlata, C., Haardt, F., et al. 2015, *ArXiv e-prints*, arXiv:1511.01998
- Sabbi, E. 2009, *Instrument Science Report WFC3 2009-19* (STScI, Baltimore)
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Sandberg, A., Östlin, G., Melinder, J., Bik, A., & Guaita, L. 2015, *ArXiv e-prints*, arXiv:1511.00684
- Schroeder, J., Mesinger, A., & Haiman, Z. 2012, *MNRAS*, 428, 3058
- Scott, J. E., Kriss, G. A., Brotherton, M., et al. 2004, *ApJ*, 615, 135
- Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
- Shapley, A. E., Steidel, C. C., Pettini, M., Adelberger, K. L., & Erb, D. K. 2006, *ApJ*, 651, 688
- Shull, J. M., Stevans, M., & Danforth, C. W. 2012, *ApJ*, 752, 162
- Siana, B., Teplitz, H. I., Colbert, J., et al. 2007, *ApJ*, 668, 62

Siana, B., Teplitz, H. I., Ferguson, H. C., et al. 2010, *ApJ*, 723, 241
 Siana, B., Shapley, A. E., Kulas, K. R., et al. 2015, *ApJ*, 804, 17
 Silk, J. 2005, *MNRAS*, 364, 1337
 Silk, J., & Norman, C. 2009, *ApJ*, 700, 262
 Silverman, J. D., Green, P. J., Barkhouse, W. A., et al. 2008, *ApJ*, 679, 118
 Silverman, J. D., Mainieri, V., Salvato, M., et al. 2010, *ApJS*, 191, 124
 Springel, V., Di Matteo, T., & Hernquist, L. 2005, *MNRAS*, 361, 776
 Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, *ApJ*, 546, 665
 Sternberg, A., McKee, C. F., & Wolfire, M. G. 2002, *ApJS*, 143, 419
 Szokoly, G. P., Bergeron, J., Hasinger, G., et al. 2004, *ApJS*, 155, 271
 Telfer, R. C., Zheng, W., Kriss, G. A., & Davidsen, A. F. 2002, *ApJ*, 565, 773
 Tumlinson, J., Giroux, M. L., Shull, J. M., & Stocke, J. T. 1999, *AJ*, 118, 2148
 Vanden Berk, D. E., Richards, G. T., Bauer, A., et al. 2001, *AJ*, 122, 549
 Vanzella, E., Siana, B., Cristiani, S., & Nonino, M. 2010a, *MNRAS*, 404, 1672
 Vanzella, E., Cristiani, S., Dickinson, M., et al. 2008, *A&A*, 478, 83
 Vanzella, E., Giavalisco, M., Inoue, A. K., et al. 2010b, *ApJ*, 725, 1011
 Vanzella, E., Guo, Y., Giavalisco, M., et al. 2012, *ApJ*, 751, 70
 Vanzella, E., de Barros, S., Castellano, M., et al. 2015, *A&A*, 576, A116
 Wilkins, S. M., Bunker, A. J., Lorenzoni, S., & Caruana, J. 2011, *MNRAS*, 411, 23

APPENDIX

A. CRITICAL TESTING OF OUR STACKING PROCEDURE AND LYC MEASUREMENTS

We performed various tests on our data to assess the robustness of our LyC stacking method and detections (§4). We test the impact of several sources of possible systematic effects or spurious signal and evaluate the overall level of confidence in our quoted uncertainties on the resulting measured LyC signal.

A.1. Detection Tests of the LyC Stacking and Measurements

To verify that residual astrometric offsets and/or trapped electron trails do not significantly affect our measurements, we first *randomly* rotate each of the individual sub-images by integer multiples of 90° , then repeat the stacking described in §4.1, as these systematics would be directionally dependent. In Fig. A1 and A2, we compare our HQ and combined HQ+IQ spectroscopic samples to the original LyC stacks (panels a–d) with the results of our rotation tests (panels e–h). The detected LyC signal in the rotated stacks does indeed remain significant at the $\gtrsim 4\sigma$ level (see Table A1). In the case of F225W, rotation of the images actually *improves* the SNR of the detection (by $\sim 20\%$), while the measured photometry remains within ~ 0.1 mag from that of the original (unrotated) stack. Such an improvement in SNR without significantly affecting the detected flux is most likely due to small-scale residual gradients in the background of the individual sub-images, which after the random rotations would average out in a stack. For the F435W filter, this is not the case, and the SNR actually decreased (by $\sim 30\%$) upon random rotation, suggesting that randomizing the surrounding sky may slightly dilute the flux, as small-scale fluctuations in the brighter background of this filter may combine with the central flux in the image (see Table A1). The F225W ERS images were all taken at the end of each orbit — closest to the Earth’s limb — since the F225W

Windhorst, R., Mathis, D., & Neuschaefer, L. 1990, in *Astronomical Society of the Pacific Conference Series (Bookcrafters, Inc.; Provo UT)*, Vol. 10, *Evolution of the Universe of Galaxies*, ed. R. G. Kron, 389
 Windhorst, R. A., & Cohen, S. H. 2010, in *American Institute of Physics Conference Series*, Vol. 1294, *American Institute of Physics Conference Series*, ed. D. J. Whalen, V. Bromm, & N. Yoshida, 225
 Windhorst, R. A., Keel, W. C., & Pascarelle, S. M. 1998, *ApJL*, 494, L27
 Windhorst, R. A., Cohen, S. H., Hathi, N. P., et al. 2011, *ApJS*, 193, 27
 Wise, J. H., & Cen, R. 2009, *ApJ*, 693, 984
 Wise, J. H., Demchenko, V. G., Halicek, M. T., et al. 2014, *MNRAS*, 442, 2560
 Wolf, C., Meisenheimer, K., Kleinheinrich, M., et al. 2004, *A&A*, 421, 913
 Worsec, G., Prochaska, J. X., Hennawi, J. F., & McQuinn, M. 2014, *ArXiv e-prints*, arXiv:1405.7405
 Wuyts, S., van Dokkum, P. G., Franx, M., et al. 2009, *ApJ*, 706, 885
 Wyithe, J. S. B., Hopkins, A. M., Kistler, M. D., Yüksel, H., & Beacom, J. F. 2010, *MNRAS*, 401, 2561
 Xue, Y. Q., Luo, B., Brandt, W. N., et al. 2011, *ApJS*, 195, 10
 Yajima, H., Choi, J.-H., & Nagamine, K. 2011, *MNRAS*, 412, 411
 Zheng, W., Kriss, G. A., Telfer, R. C., Grimes, J. P., & Davidsen, A. F. 1997, *ApJ*, 475, 469
 Zurita, A., Beckman, J. E., Rozas, M., & Ryder, S. 2002, *A&A*, 386, 801

sky-background was expected to be the faintest. Hence, slight amounts of Earth-shine may have contributed to somewhat larger sky-gradients in the F225W mosaics than in the other filters.

Next, we perform a split-halves test, where we randomly select a subset of half of the sub-images used in each stack and stack these. The remaining second half of the sub-images are similarly combined into a comparison stack. The resulting stacks are shown in Fig. A1 for the HQ sample (panels i–l and m–p) and in Fig. A2 for the combined HQ+IQ samples (same panel layout as in Fig. A1). The flux measurements resulting from the split-halves tests for the HQ+IQ galaxy samples are listed in Table A1. Each stack for each subset yields LyC fluxes at the $\gtrsim 4\sigma$ level — i.e., a significance level similar to the original stacks. It is possible that the SNR of the resulting LyC stacks depend on the particular subset of galaxy sub-images included in each sub-stack. However, the samples in each of these sub-sets are too small for a more detailed analysis of such an effect. Table A1 does show that the average LyC detected in each of the sub-sets is — to within the errors — consistent with that of the total stack (for details, see §A.2).

Last, we also extract from the WFC3/UVIS ERS mosaics random patches of *blank sky*, equal in number to the number of galaxy sub-images used for the stacks for each filter. These are combined using the same stacking method, and have the *same* sensitivity to LyC emission as the galaxy stacks. We present the resulting blank sky stacks in the bottom row (panels q–t) of Fig. A1 and A2, and their blank-sky “photometry” is tabulated in Table A1. No significant signal is detected at ≥ 31.5 mag (i.e., the central aperture “flux” is present at the $\sigma < 1$ level) in any of the blank sky stacks, which implies that our more significantly detected LyC signal in the other cases discussed above is real, and associated with objects that were selected at longer (UVC) wavelengths.

A.2. SEXTRACTOR versus “Tic-tac-toe” Photometry

It may be of some concern that our dependence on SEXTRACTOR for photometric measurements might cause us to miss some amount of extended and faint LyC flux at larger




Figure AA1: Tests for systematics in our stacking procedure for the four different filters (redshift bins) for all galaxies with HQ spectra. For reference, we repeat in panels [a] through [d] the corresponding panels from Fig. 6. In panels [e] through [h] we present the result of our test with all images rotated over random multiples of 90° before stacking; in panels [i] through [l] we show the result of randomly dividing our sample in two and stacking the first independent subset of images; in panels [m] through [p] we show the complementary stack for the remainder of the images; and in panels [q] through [t] we show that no excess LyC signal is detected in stacks of equal numbers of *random* blank sky areas. The meaning of blue and green ellipses is as in Fig. 6–8 and Fig. B1–B3.

galactocentric radii, and that its fitted measurement apertures may be smaller than desired.

We therefore analyzed the flux and SNR within the central 24×24 pixels ($2''.16 \times 2''.16$) of a “tic-tac-toe” 9-segment grid with respect to the background level, corrected for residual gradients, determined from the eight surrounding segments. An example is shown in Fig. A3. The final columns of Table A1 compare the SNR measured using our “tic-tac-toe” photometry and those measured within the SEXTRACTOR LyC apertures. We measure similar fluxes — at slightly lower SNR — using the “tic-tac-toe” aperture compared to

the smaller SEXTRACTOR fitted apertures and measurements, validating their robustness against modest variations in choice of aperture size, and the specific details of the sky-background subtraction.

The details of our “tic-tac-toe” photometry results are shown in Table A1, which serves to verify the quantities listed in Table 2 for the LyC and UVC apertures used for LyC detections in §4.2. Col. 1–3 of Table A1 are the same as Col. 1, 2 & 4 of Table 2; Col. 4 lists the measured ERS sky-background *before* the sky-level itself was removed from each exposure in the AstroDrizzle reduction (for details, see




Figure AA2: As Fig. A1 for all galaxies with HQ+IQ spectra. For reference, we repeat in panels [a] through [d] the corresponding panels from Fig. B1.

W11 Table 2); Col. 5–6 lists the sky-subtracted LyC and UVC fluxes and their formal 1σ errors from the “tic-tac-toe” photometry; Col. 7–8 lists the resulting total LyC and UVC AB-magnitudes measured over the full $2''.16 \times 2''.16$ “tic-tac-toe” apertures, which are to be compared with the same quantities derived from the SExtractor apertures in Col. 6–8 and 10–14 of Table 2; Col. 9 of Table A1 lists the corresponding *average* LyC and UVC SB-values inside the $2''.16 \times 2''.16$ apertures; Col. 10 lists the 1σ SB-error on this value *implied* by the observed total SNR of the LyC or UVC flux, assuming a fully flat SB-distribution inside the central “tic-tac-toe” aperture (see §5.4); Col. 11 shows the SNR predicted by the WFC3/UVIS and ACS/WFC CCD equation as a check of the observed SNR derived from the “tic-tac-toe” analysis, which is listed in Col. 12.

When averaged over the 28 cases listed in Table A1, the SNR predicted by the CCD detector properties is about 0.87 ± 0.30 (rms) $\times$ the observed SNR that was derived from the “tic-tac-toe” photometry. This is likely because the SNR predicted from the CCD equation has an uncertain component related to how detector read-noise and dark-current noise affect the drizzled images in a *correlated* fashion. These noise components are only measured in the original CCD images (see W11 and Dressel et al. (2015) for a discussion of all WFC3 CCD parameters). For the CCD equation prediction of the drizzled mosaics, we assumed that this term would scale with the original/drizzled pixel ratio. Depending on how large a fraction of the read-noise and dark-current is correlated on scales of a few pixels, this may not be exactly true. In any case, with the above assumption, the predicted

SNR for the $2''.16 \times 2''.16$ “tic-tac-toe” apertures is close to the SNR derived from the observed parameters in the “tic-tac-toe” photometry. More specifically, the observed “tic-tac-toe” SNR values in Col. 12 of Table A1 are generally close to the SNR derived from the SExtractor photometry that was measured over the smaller LyC and UVC apertures (see §4.2 and Table 2). The exception is the F435W photometry, where the “tic-tac-toe” SNR appears to be lower for the HQ sample. This is likely due to the larger amount of higher sky-background and its sky-background gradients included in the “tic-tac-toe” aperture in the ACS/WFC F435W filter, compared to the WFC3 UVIS filters.

Table 1 lists the measured 1σ SB-sensitivity over a $2''.0$ diameter aperture in our drizzled WFC3 and ACS mosaics. For example, the F275W filter has a single mosaic 1σ SB-sensitivity of $\sim 29.82 \text{ mag arcsec}^{-2}$ when measured over a $2''.0$ diameter aperture. If the sky-background was completely flat without any gradients (but see the discussion in Appendix C.1) then in a 26-object HQ+IQ galaxy stack with 8 surrounding “tic-tac-toe” apertures, the quality of the sky-subtraction, or the 1σ SB-sensitivity, could in principle be as good as $29.82 \text{ mag} + 2.5 \log \sqrt{(2.16/2)^2 \times 26 \times 8} = 32.80 \text{ mag arcsec}^{-2}$. Here, we scaled the 1σ SB-sensitivity with the aperture size, and the rms deviation of the surrounding sky with the number of sub-images and number of available surrounding sky-boxes (each of which have their own well measured residual sky-values and sky-rms; see Fig. A3). All of these values are determined *after* image stacking, and are used in predicting the “tic-tac-toe” SNR in Table A1.

This limit is marginally brighter than the measured 1σ SB-sensitivity value derived from the “tic-tac-toe” photometry in Table A1 ($33.60 \text{ mag arcsec}^{-2}$), which could be due to some correlated or systematic noise present in the actual data. Nevertheless, since subtle residual gradients remain in the sky-background, as discussed in Appendix C.1, the real 1σ SB-sensitivity limit that can be achieved is *not as faint* as either one of these values. We suggest in Appendix C.1 that uncertainties from subtle residual sky-background gradients in the LyC stacks limit our photometry to $\sim 32.0-32.5 \text{ mag arcsec}^{-2}$, so that any LyC SB values measured to be fainter than this should be considered with this limit in mind. For that reason, we do not plot any of the SB-values in the LyC light-profiles of Fig. 10 to levels fainter than $32.0 \text{ mag arcsec}^{-2}$. Also, when the formal “tic-tac-toe” 1σ SB-sensitivity (Col. 10 of Table A1) is much fainter than the limit of $\sim 32.0-32.5 \text{ mag arcsec}^{-2}$ from residual sky-background systematics (Appendix C.1), the SNR derived for the larger “tic-tac-toe” apertures may be overestimated. Hence, for $SB \gtrsim 32.0-32.5 \text{ mag arcsec}^{-2}$ in Col. 10 of Table A1, the quoted SNR values must be interpreted accordingly. As quantified in Appendix A.3, the “tic-tac-toe” photometry does confirm the SExtractor photometry from the smaller LyC and UVC apertures discussed in §4.2, to within their respective errors.

A.3. Quality of SExtractor versus “Tic-tac-toe” Photometry

When compared to our main SExtractor photometry in Table 2, the “tic-tac-toe” photometry in Table A1 shows the following: for galaxies with HQ spectra, the measured difference in flux within the LyC detected apertures and the $2''.16 \times 2''.16$ “tic-tac-toe” aperture is $\Delta(\text{LyC-TTT}) = +0.14 \pm 0.21 \text{ mag (m.e.)}$, when averaged over the four LyC filters. For the galaxies with HQ+IQ spectra, this difference is $\Delta(\text{LyC-TTT}) = +0.30 \pm 0.09 \text{ mag (m.e.)}$. For the galaxies with HQ spectra, this flux difference is not significant, while for the HQ+IQ sample, the difference is significant, and indeed goes in the direction of the larger “tic-tac-toe” apertures containing slightly more LyC flux. This is possibly another sign of the very flat LyC SB-profiles discussed in §5.4.

In the “tic-tac-toe” exercise, all contaminating neighbors were removed from the central $2''.16 \times 2''.16$ “tic-tac-toe” aperture. Nonetheless, it is also possible that some contaminating very low-SB flux (invisible in the individual sub-images, even in our deepest WFC3 IR-images) from nearby neighbors could have leaked into the larger “tic-tac-toe” apertures. In any case, the above numbers show that the total contaminating flux is likely $\lesssim 0.3 \text{ mag}$, since any contaminating flux should be far smaller in the much smaller LyC apertures (see also §4.2).

The “tic-tac-toe” photometry in Table A1 also allows us to further quantify the three other critical tests that were qualitatively discussed in Appendix A.1. For the stack of all galaxies with HQ and IQ spectra, the “tic-tac-toe” photometry of the sub-images that were randomly rotated by 90° shows a difference with the “unrotated” image stacks of $\Delta(\text{Rotated-Unrotated}) = +0.05 \pm 0.06 \text{ mag (m.e.)}$ when averaged over the four LyC filters. In other words, the “tic-tac-toe” photometry of the rotated HQ+IQ galaxy stack is consistent with that of the corresponding unrotated stack, showing that — to within the errors — our stacking method yields reproducible and consistent LyC fluxes.

For the sample of all galaxies with HQ+IQ spectra, the “tic-tac-toe” photometry of the first and second randomly selected sub-image halves show a difference with the unrotated stack of $\Delta(1\text{st half}-2\text{nd half}) = +0.17 \pm 0.19 \text{ mag (m.e.)}$ when averaged over the four LyC filters. This is not a significant difference, so the “tic-tac-toe” photometry on the two independent sub-halves of the data gives to within $\lesssim 0.2 \text{ mag}$ the same total LyC flux as the total sample. The noticeable exception is again the ACS/WFC F435W image stack, where the sky-level and residual sky-gradients in the “tic-tac-toe” apertures are greater, resulting in an overall larger uncertainty in the measured LyC fluxes in F435W.

No signal was detected (at the $AB \lesssim 31.5 \text{ mag}$ level) in any of the random blank-sky central “tic-tac-toe” apertures, when compared to the average surrounding sky-boxes. These “blank-sky” stacks thus serve as a check on the quality of the UV sky-background subtraction, and the effects of any subtle remaining sky-gradients. They also show that in random sky-areas, no significant amount of flux from cosmic rays residuals that might have not been completely removed during the drizzling process were added in at the $AB \lesssim 31.5 \text{ mag}$




Figure AA3: Example of a “tic-tac-toe” sky-background analysis of the 71×71 pixel LyC [left 4 panels] and UVC [right 4 panels] sub-image stacks. Any large-scale gradients in the residual sky-background left in the drizzled images are $\gtrsim 5.2\text{--}40\times$ fainter than the faint remaining sky background residuals determined in Fig. 3. Such gradients in the sky will be fainter than $\sim 32.1, 32.0, 32.4$, and 32.0 mag arcsec $^{-2}$ across the $6''.39$ “tic-tac-toe” aperture in the WFC3/UVIS F225W, F275W, and F336W, and ACS/WFC F435W sub-image stacks. Residual UV sky-gradients appear, therefore, fainter than the LyC signal where this can be measured, as discussed in the text. Not all “tic-tac-toes” used in Table A1 are shown here.

level (see Fig. A1–A2). Hence, these blank-sky stacks are our best check that even for $N=3\text{--}6$ one-third to half-orbit exposures per filter, the bulk of our stacked, very faint LyC signal at $m_{\text{AB}} \sim 29.5\text{--}30.7$ mag (Tables 2 and A1) is not due to residual unfiltered cosmic rays or noise peaks. If this were true, then these random blank-sky stacks would have shown as significant false signal at *similar* m_{AB} -levels as the real LyC detections, which was not the case.

B. POSSIBLE SOURCES OF CONTAMINATING NON-IONIZING FLUX

B.1. In-filter Red-leak of Non-ionizing Flux

The WFC3/UVIS and ACS/WFC filters were designed to minimize the transmission of photons with wavelengths higher or lower than their specified cutoffs (see Fig. 1(a)). However, as seen in Fig. 1(b), a small amount of flux redward of the Lyman Limit from galaxies observed in these filters with redshifts in the ranges of Table 1 can still leak into the filter and contaminate LyC observations with non-ionizing UVC flux. The lower redshift-bounds in Table 1 were carefully chosen such that *no* light $> 912\text{\AA}$ is sampled below the filter’s red-edge. The filter red-edge is defined as $< 0.5\%$ of the filter’s peak transmission. For galaxies at the higher redshifts in the ranges of Table 1, and especially those at higher redshifts than the designated upper bound, the contribution from UVC “red-leak” can become the dominant source of photons measured in the filter, as the portion of the spectrum intended for LyC observation becomes exceedingly faint at shorter wavelengths and the non-ionizing continuum remains roughly constant at longer wavelengths. Thus, in order to ac-

curately measure LyC photometry and escape fractions from these samples of galaxies, we must verify that the flux measured from our sample is dominated by LyC photons.

Since we cannot directly measure the fraction of non-ionizing flux leaking into the filter from the observation, we estimate this value by modeling the contribution of LyC and UVC to the observed flux from the total sample. Using SEDs fit from continuum observations of our galaxy sample and *average* line-of-sight IGM transmission models (see §5.1), we calculate the average UVC “red-leak” of our observation in the WFC3/UVIS F225W, F275W, F336W, and ACS/WFC3 F435W filters by comparing the total flux integrated in the entire filter, and the total flux integrated below the Lyman Limit of each galaxy. We calculate this value as:

$$\frac{F_{\nu}^{\text{UVC}}}{F_{\nu}^{\text{LyC}}} = 1 - \frac{\sum_{i=1}^{N_{\text{gal}}} \int_{\lambda_1}^{\lambda_{912}} \mathcal{T}_{\text{obs}}^{\text{LyC}}(\nu) \mathcal{T}_{\text{IGM}}(z_i, \nu) F_{\nu,i}(\nu) \frac{d\nu}{\nu}}{\int_{\lambda_1}^{\lambda_{912}} \mathcal{T}_{\text{obs}}^{\text{LyC}}(\nu) \frac{d\nu}{\nu}} \bigg/ \frac{\int_{\lambda_1}^{\lambda_2} \mathcal{T}_{\text{obs}}^{\text{LyC}}(\nu) \mathcal{T}_{\text{IGM}}(z_i, \nu) F_{\nu,i}(\nu) \frac{d\nu}{\nu}}{\int_{\lambda_1}^{\lambda_2} \mathcal{T}_{\text{obs}}^{\text{LyC}}(\nu) \frac{d\nu}{\nu}}$$

where λ_1 and λ_2 are the minimum and maximum wavelengths of the full filter transmission curve, λ_{912} is the observed wavelength of the Lyman Limit of the galaxy, F_{ν} is the SED flux of the galaxy in $\text{erg s}^{-1} \text{cm}^{-2} \text{Hz}^{-1}$, $\mathcal{T}_{\text{IGM}}$ is the average line-of-sight IGM transmission at the redshift of the galaxy, and $\mathcal{T}_{\text{obs}}^{\text{LyC}}(\nu)$ is the combined throughput of the filter, detector quantum efficiency (QE), and optical telescope assembly (OTA). This value quantifies the fraction of flux we measure from these galaxies in the filter intended for LyC observations that is non-ionizing. For the F225W, F275W, F336W, and F435W filters, the *percentage of total* “red-leak” photons

Table AA1.: Photometric “Tic-Tac-Toe” Analysis of the LyC and UVC from Galaxies with HQ and HQ+IQ Spectra, and Other Photometry Tests

Filter	z -range	N_{objects}	μ_{sky} [mag r''^2]	$S_{\text{LyC/UVC}}$ [$e^-/\text{pix/s}$]	$\sigma_{S_{\text{LyC/UVC}}}$ [$e^-/\text{pix/s}$]	$S_{\text{LyC/UVC}}$ [m_{AB}]	σ_S [m_{AB}]	$\mu_{\text{LyC/UVC}}$ [mag r''^2]	$1-\sigma_\mu$ [mag r''^2]	SNR (Pred.)	SNR (Obs.)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
LYC - GALAXIES WITH AGN (HQ SPECTRA):											
F225W	2.302–2.450	1	25.46	4.73×10^{-5}	8.95×10^{-6}	27.95	0.21	29.63	31.43	3.14	5.28
F275W	2.559–3.076	7	25.64	2.12×10^{-5}	3.43×10^{-6}	28.91	0.18	30.58	32.56	4.32	6.20
F336W	3.132–3.917	4	24.82	2.10×10^{-5}	4.87×10^{-6}	29.46	0.25	31.13	32.72	3.51	4.31
F435W	4.414–5.786	2	23.66	3.39×10^{-6}	7.11×10^{-5}	>29.1	—	>30.8	30.82	(1 σ)	—
UVC - GALAXIES WITH AGN (HQ SPECTRA):											
F606W	2.302–2.450	1	22.86	3.87×10^{-3}	2.42×10^{-4}	25.62	0.07	27.29	30.30	15.0	16.0
F606W	2.559–3.076	7	22.86	7.74×10^{-3}	5.45×10^{-5}	24.86	.007	26.54	31.92	71.6	141.
F775W	3.132–3.917	4	22.64	6.19×10^{-3}	1.55×10^{-4}	24.27	0.03	25.95	29.95	44.1	39.9
F850LP	4.414–5.786	2	22.58	2.52×10^{-3}	7.17×10^{-5}	24.46	0.03	26.13	29.99	28.4	35.1
LYC - GALAXIES WITHOUT AGN (HQ SPECTRA):											
F225W	2.302–2.450	14	25.46	8.98×10^{-6}	1.52×10^{-6}	29.76	0.18	31.43	33.36	3.64	5.90
F275W	2.559–3.076	11	25.64	1.12×10^{-5}	3.62×10^{-6}	29.61	0.35	31.28	32.50	3.58	3.08
F336W	3.132–3.917	10	24.82	8.37×10^{-6}	3.36×10^{-6}	30.46	0.44	32.13	33.12	2.22	2.49
F435W	4.414–5.786	15	23.66	1.49×10^{-5}	1.17×10^{-5}	30.84	0.86	32.51	32.78	1.16	1.26
UVC - GALAXIES WITHOUT AGN (HQ SPECTRA):											
F606W	2.302–2.450	14	22.86	1.24×10^{-2}	7.87×10^{-5}	24.36	.007	26.03	31.52	156.	157.
F606W	2.559–3.076	11	22.86	1.16×10^{-2}	6.89×10^{-5}	24.42	.006	26.10	31.66	125.	168.
F775W	3.132–3.917	10	22.64	3.66×10^{-3}	3.02×10^{-5}	24.85	.009	26.52	31.73	60.1	121.
F850LP	4.414–5.786	15	22.58	6.77×10^{-4}	2.31×10^{-5}	25.89	0.04	27.56	31.22	37.3	29.3
LYC - GALAXIES WITHOUT AGN (HQ+IQ SPECTRA):											
F225W	2.262–2.450	29	25.46	1.05×10^{-5}	9.94×10^{-7}	29.59	0.10	31.26	33.82	8.60	10.6
F275W	2.481–3.076	26	25.64	1.45×10^{-5}	1.32×10^{-6}	29.33	0.10	31.00	33.60	10.6	10.9
F336W	3.110–4.149	23	24.82	1.69×10^{-5}	2.14×10^{-6}	29.70	0.14	31.37	33.62	5.57	7.91
F435W	4.414–6.277	37	23.66	4.22×10^{-5}	7.09×10^{-6}	29.71	0.18	31.38	33.32	7.85	5.96
UVC - GALAXIES WITHOUT AGN (HQ+IQ SPECTRA):											
F606W	2.262–2.450	29	22.86	1.13×10^{-2}	4.09×10^{-5}	24.46	.004	26.13	32.23	237.	275.
F606W	2.481–3.076	26	22.86	9.12×10^{-3}	2.52×10^{-5}	24.69	.003	26.36	32.75	190.	362.
F775W	3.110–4.149	23	22.64	4.30×10^{-3}	4.11×10^{-5}	24.67	0.01	26.34	31.39	122.	104.
F850LP	4.414–6.277	37	22.58	4.22×10^{-4}	8.08×10^{-6}	26.40	0.02	28.07	32.36	54.9	52.2
LYC - GALAXIES WITHOUT AGN (HQ+IQ SPECTRA) — ROTATED BY RANDOM $N \times 90^\circ$:											
F225W	2.262–2.450	29	25.46	1.08×10^{-5}	7.83×10^{-7}	29.56	0.08	31.23	34.08	8.84	13.7
F275W	2.481–3.076	26	25.64	1.47×10^{-5}	1.35×10^{-6}	29.31	0.10	30.98	33.58	10.8	10.9
F336W	3.110–4.149	23	24.82	1.64×10^{-5}	2.06×10^{-6}	29.73	0.14	31.40	33.65	5.40	7.94
F435W	4.414–6.277	37	23.66	3.46×10^{-5}	7.67×10^{-6}	29.92	0.24	31.60	33.23	6.48	4.51
LYC - GALAXIES WITHOUT AGN (HQ+IQ SPECTRA) — FIRST RANDOM HALF OF ALL SUB-IMAGES:											
F225W	2.262–2.450	14	25.46	1.20×10^{-5}	2.02×10^{-6}	29.44	0.18	31.11	33.05	4.85	5.95
F275W	2.481–3.076	13	25.64	1.18×10^{-5}	1.22×10^{-6}	29.55	0.11	31.22	33.69	4.47	9.72
F336W	3.110–4.149	11	24.82	1.28×10^{-5}	3.10×10^{-6}	30.00	0.26	31.67	33.21	2.06	4.14
F435W	4.414–6.277	18	23.66	9.45×10^{-5}	1.96×10^{-5}	28.83	0.22	30.51	32.22	8.51	4.83
LYC - GALAXIES WITHOUT AGN (HQ+IQ SPECTRA) — SECOND RANDOM HALF OF ALL SUB-IMAGES:											
F225W	2.262–2.450	15	25.46	1.12×10^{-5}	1.03×10^{-6}	29.51	0.10	31.19	33.78	4.85	10.9
F275W	2.481–3.076	13	25.64	1.76×10^{-5}	1.88×10^{-6}	29.12	0.12	30.79	33.22	6.59	9.38
F336W	3.110–4.149	12	24.82	2.12×10^{-5}	1.73×10^{-6}	29.45	0.09	31.12	33.85	3.68	12.3
F435W	4.414–6.277	19	23.66	7.60×10^{-5}	1.65×10^{-5}	29.07	0.24	30.74	32.40	7.27	4.61
RANDOM EMPTY SKY:											
F225W	2.262–2.450	29	25.46	5.44×10^{-7}	1.52×10^{-6}	>31.7	—	>33.3	33.35	(1 σ)	—
F275W	2.481–3.076	26	25.64	3.29×10^{-7}	1.30×10^{-6}	>31.9	—	>33.6	33.61	(1 σ)	—
F336W	3.110–4.149	23	24.82	7.30×10^{-8}	9.71×10^{-7}	>31.6	—	>33.3	33.26	(1 σ)	—
F435W	4.414–6.277	37	23.66	1.02×10^{-6}	1.95×10^{-5}	>30.5	—	>32.2	32.22	(1 σ)	—

(1) WFC3 or ACS filter; (2) redshift range (as in Table 2); (3) Number of galaxies with high or intermediate quality spectroscopic redshifts used in each stack; (4) sky surface brightness in AB mag arcsec $^{-2}$ using EXPTIME and MDRZSKY from the FITS header, corrected for the number of tiles in the mosaic; (5)–(6) Average sky-subtracted flux and error thereon in $e^- \text{ pix}^{-1} \text{ s}^{-1}$ over N sub-images in the central 24×24 pixel “tic-tac-toe” aperture (see Appendix A.2); (7)–(8) Total LyC and UVC flux and error thereon, expressed as AB mag; (9)–(10) Average LyC and UVC surface brightness and 1σ error thereon, both in AB mag arcsec $^{-2}$; (11) SNR of LyC or UVC flux in Col. (5), predicted from the WFC3/UVIS or ACS/WFC CCD equation; (12) SNR of LyC or UVC detection in Col. (5), observed from the total sky-subtracted LyC (row 1–4) or UVC (row 5–8) flux and corresponding sky subtraction error in Col. (6). (These are to be compared with the SNR of LyC or UVC detections within the SExtractor LyC or UVC apertures listed in Table 2 or Table B1).

that contribute to the measured LyC flux of our sample are $\sim 0.65\%$, 0.64% , 0.19% , and 0.66% , respectively. That is, less than 1% of the anticipated and measured LyC flux itself could be redleak flux from longwards of $912\text{\AA}$. From these values, it is clear that LyC observations of galaxies at the redshift ranges indicated in Table 1 with their respective filters are dominated by LyC photons. We also note that our MC analysis of the observed LyC flux from these galaxies accounts for these “red-leak” photons, in order to make appropriate corrections for low-level non-ionizing contamination of the order of $\sim 0.28\%$.

B.2. UVIS-filter Pinholes

Pinholes are very small voids in the coating on the surface of a filter. These voids appear usually due to poor adhesion of the coating in these regions where particulate matter on the surface of the filter is coated-over when the substrate is cast, or from mechanical abrasion or chemical interactions when the filter is in use. Several of the WFC3/UVIS filters have pinholes, so we must make sure that none of the LyC flux that we measure is due to out-of-band flux leaking in through the filter in an area where such a pinhole exists. Most of the obvious pinholes were known before WFC3’s launch, and the filters with the fewest pinholes were chosen for flight (Dressel et al. 2015). To the best of our knowledge, the number of pinholes did not increase during the 7 years that the WFC3 filters were on the ground. Visible pinholes on the selected filters were painted over when possible (Baggett et al. 2006). Any remaining pinholes not painted over before launch are likely $\lesssim 0.2$ mm in diameter, or they would have been treated before final instrument assembly. Unfortunately, no record was kept of any less obvious pinholes in the flight filters that were not painted over before launch. Remaining pinholes could cause subtle field-dependent redleaks and very low-level sky-gradients, which we quantify here.

We need to first estimate how large the footprint and the amplitude of any pinhole redleak on the WFC3 CCDs could be. The HST f/24 beam gets re-imaged inside WFC3 to f/31 (Dressel et al. 2015), so that the plate-scale on the WFC3 UVIS detector changes from $3''.58/\text{mm}$ to $2''.77/\text{mm}$ (i.e., $206265/(2400 \times 31)''/\text{mm}$). The WFC3 UVIS Marconi CCDs have $15\mu\text{m}$ pixels, so the two $2\text{k} \times 4\text{k}$ CCD-arrays are about 61 mm in physical size. The WFC3/UVIS F225W filter is in Selectable Optical Filter Assembly (SOFA) filter wheel 3, F336W in filter wheel 4, and F275W in filter wheel 6 out of 12, where wheel 12 is closest to the CCD’s. The average location of these three filter wheels is about 190 ± 25 mm away from the focal plane (Fig. 2.1 of Dressel et al. 2015), which we hereafter refer to as the “center of the SOFA”. The ± 25 mm indicates the approximate range over which these three UVIS filter wheels are mounted inside the SOFA. Each SOFA filter is 57.3 mm square and $\lesssim 5.151$ mm thick (Baggett et al. 2006), as fabricated by the filter vendor to the specifications required by the WFC3 Scientific Oversight Committee and Instrument Product Team.

The SOFA is about 1/3 of the way between the focal plane and the pupil, which is the anamorphic asphere mirror inside WFC3 that corrects for the spherical aberration in HST’s

primary mirror. Fortunately therefore, all pinholes in the WFC3 UVIS filters will be severely out of focus, since the filters are so far from the focal plane. We first need to calculate how large the pupil of each image is in the filter plane. The anamorphic asphere mirror has a diameter of about 25 mm and is about 630 mm away from the CCD. It is about 440 mm from the center of the SOFA, so that the *radius* of the image pupil at the filter distance is about $r_{\text{ip-f}} \simeq (190/630) \cdot 25/2 \simeq 3.77$ mm. Hence, the image pupil at the filter is about $7.54/57.3 \simeq 13\%$ of the filter size.

Next, we need to estimate how large the footprint and the amplitude of any pinhole flux on the WFC3 CCDs could be. If WFC3’s f/31 beam goes through a pinhole with an $r \gtrsim 0.08$ mm radius (i.e., $\gtrsim 0.5 \times 5.151/31$) in a $\lesssim 5.151$ mm thick UV filter about 190 mm in front of the CCD, this pinhole will affect a beam with an opening angle $\theta = 90^\circ - \text{atan}(31) \simeq 1.85^\circ$ projected onto the CCD. As viewed from the CCD, the remainder of the pixels outside this beam will not view the sky through the pinhole. At the CCD, the circular beam that is affected by this pinhole will have a $190/31 \simeq 6.1$ mm $\simeq 17''.0 \simeq 430$ pixel radius on the CCD, and so its diameter will cover about 20% of the WFC3 CCD-FOV. To avoid internal reflections in the camera, the UVIS CCDs are tilted by $\sim 21^\circ$ with respect to the axis of the beam, so the projected footprint of each pinhole is actually about $430/\cos(21) \simeq 460$ pixels in radius. In other words, the footprint projected by the pinhole on the CCD is very large, and will significantly dilute the extra SB signal projected through the pinhole. In the limit, a much smaller pinhole ($r < 0.08$) mm would not see the entire f/31 beam, and will thus act like a pinhole camera that illuminates the entire CCD, diluting the extra SB that goes through this smaller pinhole even more.

If the interference or AR coating were not present in a pinhole for one of our WFC UVIS filters, Fig. 1b shows that it could have a significant increase in local throughput, or less in case that the local defect was only partial in transmission. In a worst case, the pinhole would act like a F606W or F775W WFC3 filter at that location, if the $\text{OTA} \times \text{Filter-Throughput} \times \text{CCD-QE}$ at those wavelengths reached the WFC3 maximum in the F606W filter of $\sim 28\%$ (see Fig. 3.2 and 5.2 of Dressel et al. 2015). In such cases, the Zodiacal sky in the beam illuminated by the pinhole would be much higher than seen in our UVIS filters, possibly as high as that in our broad-band optical filters, or slightly higher if the pinhole were fully transparent.

The brightest object in the WFC3 ERS has $V \simeq 17 m_{\text{AB}}$ and the most commonly seen objects are faint galaxies with $V \simeq 26\text{--}27 m_{\text{AB}}$ (Fig. 12 of W11). Hence, their collective SB is well below the average Zodiacal background, which is $23.7\text{--}22.6$ mag arcsec $^{-2}$ (see Col. 4 of Table A1). Hence, for all practical purposes, the pinhole contribution will just be the full white-light Zodiacal background if its throughput reaches the maximum total throughput of $\sim 28\%$. Following Table A1, we assume that the white-light Zodi would have on average a $\text{SB} \simeq 22.9$ mag arcsec $^{-2}$ through such a pinhole, which is roughly the observed value in F606W (see W11). We will also consider the case of a single $V \simeq 17 m_{\text{AB}}$ star shining behind a pinhole, as well as the integrated

sky SB derived from the faint galaxy counts in Fig. 12 of W11 to $m_{AB} \lesssim 26$ mag. In F606W, the latter reaches 10^5 galaxies/ 0.5 mag/deg^2 to $m_{AB} \lesssim 26$ mag, and in F275W, they reach 3×10^4 galaxies/ 0.5 mag/deg^2 to $m_{AB} \lesssim 25.5$ mag. On a per square arcsecond basis, the integrated sky-SB from faint galaxies is therefore $\sim 29.3 \text{ mag arcsec}^{-2}$ in F606W and $\sim 30.1 \text{ mag arcsec}^{-2}$ in F275W, respectively, i.e., fully negligible compared to the Zodiacal sky-SB values from Table A1 of $SB_V^{\text{Zodi}} \simeq 22.9$ and $SB_{\text{NUV}}^{\text{Zodi}} \simeq 25.5 \text{ mag arcsec}^{-2}$ in these filters, respectively. Hence, only the Zodiacal light and the effective SB of the occasional bright star behind the filter would be the main sources of pinhole contamination. All calculations below are done in terms of surface brightness (SB in mag arcsec^{-2}) or Intensity (I in relative counts/sec).

In a slow $f/31$ beam, when the pinhole is larger than the minimum size to transmit through the filter, the total white-light transmitted would increase proportionally to the pinhole area compared to the total area of the image pupil at the filter $(r_{\text{ph}}/r_{\text{ip-f}})^2$, both measured in mm. For an untreated pinhole with an assumed $r_{\text{ph}} \simeq 0.1$ mm, we can now estimate the increase in sky-SB contribution from this pinhole over a $r=460$ pixel radius on the CCD. In relative units, this increase is:

$$I_V^{\text{ph}} = F \cdot [r_{\text{ph}}^2/r_{\text{ip-f}}^2] \cdot [I_V^{\text{Star}} + I_V^{\text{Zodi}}] \quad (\text{B1})$$

with $I_V^{\text{Star}} = 10^{-0.4(SB_V^{\text{Star}} - ZP_V)}$, $I_V^{\text{Zodi}} = 10^{-0.4(SB_V^{\text{Zodi}} - ZP_V)}$, and:

$$I_{\text{NUV}}^{\text{ph}} = [(r_{\text{ph}}^2 - r_{\text{ip-f}}^2)/r_{\text{ip-f}}^2] \cdot [I_{\text{NUV}}^{\text{Star}} + I_{\text{NUV}}^{\text{Zodi}}] \quad (\text{B2})$$

$I_{\text{NUV}}^{\text{Star}} = 10^{-0.4(SB_{\text{NUV}}^{\text{Star}} - ZP_{\text{NUV}})}$, $I_{\text{NUV}}^{\text{Zodi}} = 10^{-0.4(SB_{\text{NUV}}^{\text{Zodi}} - ZP_{\text{NUV}})}$. Eq. B1 describes the relative counts of optical white light through the pinhole, and Eq. B2 the relative counts for the uncorrupted NUV sky-SB. Note that the optical white-light $SB_V^{\text{Zodi}} \simeq 22.9 \text{ mag arcsec}^{-2}$ is compared here to the WFC3 F606W zeropoint of $ZP_V = 26.08$ mag (i.e., *not* the ACS ZP in this case), and the original UV $SB_{\text{NUV}}^{\text{Zodi}} \simeq 25.5 \text{ mag arcsec}^{-2}$ (Col 4 of Table A1) is compared to the combined F225W and F275W zeropoints of ~ 24.1 mag. The factor $F \simeq (10000-4000)/2000 \simeq 3$ reflects that the pinhole could transmit three broadband filters worth of white light from $4000-10,000 \text{ \AA}$ (see Fig. 1b). We find approximately the same values if we instead use the WFC3 white-light filter F200LP and its zeropoint $ZP = 27.36$ mag, and set the factor $F=1$. The limit of $r_{\text{ph}} \simeq r_{\text{ip-f}}$ would describe a hypothetical pinhole so large that it transmits full white light over the entire image pupil at the filter, which now acts like a wide V -band filter. In that case, both equations still give the correct results: Eq. B1 describes its wide V -band SB, and Eq. B2 describes its now vanishingly small NUV SB.

At first, we ignore the terms with I_V^{Star} and $I_{\text{NUV}}^{\text{Star}}$ due to a bright star near a pinhole. The SB from the pinhole in V and NUV then are $I_V^{\text{ph}} \simeq 0.039$ and $I_{\text{NUV}}^{\text{ph}} \simeq 0.28$ in the same relative units, respectively. The ratio $I_V^{\text{ph}}/I_{\text{NUV}}^{\text{ph}}$ is 0.14, so that about 14% of white-light background through the pinhole gets added to the UV-sky:

$$SB_{\text{NUV}}^{\text{Sky}} = SB_{\text{NUV}}^{\text{Zodi}} - 2.5 \log(I_V^{\text{ph}}/I_{\text{NUV}}^{\text{ph}}) \quad (\text{B3})$$

The error on this is at least $25/190 \sim 0.13$ mag, depending how

far the SOFA filter is from the CCD. That is, a full white-light pinhole footprint could add $\sim 14 \pm 2\%$ to the sky-SB over a circular donut-shaped area with a diameter about 20% of the 61 mm CCD area. This is the worst case — a smaller pinhole that doesn't project the entire $f/31$ beam through the filter could instead add a much fainter SB over the whole chip.

If we also add the effect from a $m_V^{\text{Star}} \simeq 17$ mag, $m_{\text{NUV}}^{\text{Star}} \simeq 18$ mag star whose image pupil in the filter plane illuminates the pinhole, we must first correct for the fact that the light from this point source is now spread out by factor of $(r_{\text{ip-f}}/r_{\text{ph}})^2$ at the filter, so we must add $2.5 \log(3.77/0.1)^2$ to the star's point source flux of $m_{AB} \sim 17-18$ mag in Eq. B1–B2 to get the equivalent SB from the star that actually affects the pinhole, expressed in the appropriate relative units. Note that the Zodiacal background and integrated galaxy counts do not have this problem, since they are already expressed as proper SB in Eq. B1–B2. These numbers are $SB_V^{\text{Star}} \simeq 24.9$ and $SB_{\text{NUV}}^{\text{Star}} \simeq 25.8 \text{ mag arcsec}^{-2}$. Eq. B1–B2 converts these SB numbers to relative fluxes, then adds them linearly. With a $V \sim 17$ mag star, the white-light SB from the pinhole now grows from $I_V^{\text{ph}} \simeq 0.039$ to $I_V^{\text{ph}} \simeq 0.046$. This is only 17% larger than just the light from Zodiacal light alone, since the star is so much more spread out behind the filter. The NUV comparison term remains at $I_{\text{NUV}}^{\text{ph}} \simeq 0.28$. Hence, an out of focus image of a $V \sim 17$ mag star behind the pinhole adds $\sim 17 \pm 2\%$ of white-light background to the NUV sky.

To first order, both the proper NUV light and any white-light pinhole flux would get flat-fielded away during drizzling, although the pinhole regions would have a different color of the sky-background than the regular UV sky. The WFC3 UVIS Marconi CCDs have QE-curve that is fairly flat as a function of wavelength, with most QE-values between $2000-8000 \text{ \AA}$ peak around $\text{QE} \sim 70\%$ with a range of $\pm 10\%$. This peak actually lies within the F275W filter. We therefore did the experiment to flat-field a WFC3 F606W image with a high SNR flat-field taken in the F275W filter, and — owing to WFC3's flat QE-curve — this did not result in a residual gradient larger than about 10% of sky corner-to-corner across the whole CCD frame. Below we will assume a worst case of $\sim 10\%$.

We can use this result to compute the additional sky-gradient that an unpainted pinhole with $r \simeq 0.1$ mm could have caused in our UV images. When a pinhole adds about 14–17% extra flux to the regular UV Zodiacal sky over a 920 pixel diameter circle on the CCD, the residual sky-gradient that the improper flat-fielding induces is at most 0.6–0.8% of the *total* Zodiacal sky over 920 native pixels, or about 404 drizzled pixels. Across our sub-image size of 71×71 drizzled pixels, this residual sky-gradient — if present in our object sub-images — is thus less than $\sim 0.22-0.26\%$ of the local Zodiacal sky. That is, the error in the UV-sky of Eq. B3 from sky-gradients induced by the partial improper flat-fielding of any pinhole white-light is:

$$\sigma_{\text{Sky}} \simeq SB_{\text{NUV}}^{\text{Sky}} + 2.5 \log(I_V^{\text{ph}}/I_{\text{NUV}}^{\text{ph}} \cdot 0.1 \cdot 71/404) \quad (\text{B4})$$

For our average UV-sky of $25.36 \text{ mag arcsec}^{-2}$ (which is now brightened by -0.14 mag due to the extra pinhole flux), the residual sky-gradients left by pinholes in 71×71 driz-

zled pixels are thus fainter than $25.36 - 2.5 \log(0.22\%) \simeq 31.9$ mag arcsec $^{-2}$ and fainter than $\simeq 31.7$ mag arcsec $^{-2}$ if a $V \simeq 17$ mag star is also nearby the pinhole. These pinhole-induced systematics are at worst slightly brighter than those possibly caused by subtle residual gradients left at the 32.3 mag arcsec $^{-2}$ level across the CCD's due to remaining errors in the bias, dark-frames, or flat-fields, as discussed in §2.3 and Appendix C.1). Of course, the latter may affect our stacks everywhere in the CCD mosaics, while the former occur only in sporadic (although unknown) locations.

We investigated if the effects of any pinhole redleaks were in fact seen in the ERS data, since obvious pinholes would appear as large donut shaped objects in the drizzled images. No obvious defects with $\gtrsim 14$ –17% increased transmission were seen in the raw ERS data on scales of ~ 920 native pixels, although this is hard to see due to cosmic rays and the shallow depth of individual images. Partial transmission defects might exist at lower levels. As discussed in §4.1, we inspected all LyC sub-images individually, and removed the ones with suspected increased noise due to residual cosmic rays, structure in the weight-maps due to drizzled image borders, and other image defects. The objects removed from the LyC stacking all appear to have higher image rms due to the proximity of structure in the weight-maps due to drizzled image borders. No obvious enhancements in the LyC signal were seen due to the proximity of bright stars.

We find that both redleak flux from UV-filter pinholes and other subtle calibration errors may result in residual sky-gradients of order ~ 32 –32.3 mag arcsec $^{-2}$ across our stacking sky-boxes. Unless these effects can be removed through further refinement of the WFC3 calibration techniques, residual systematic subtraction errors of order $\simeq 32$ mag arcsec $^{-2}$ may well pose a fundamental limit to the LyC stacking method of WFC3 data. This is why the light-profiles in Fig. 10 cannot be extended to SB-levels much fainter than $AB \simeq 32$ mag arcsec $^{-2}$ (see also the discussion in Appendix A.2 and Table A1).

Finally, if our LyC detections were in fact caused by subtle sky-gradients on ~ 920 pixel scales, they should have been *preferentially* located near the same physical regions of the CCD, as would have been the case with CTE-effects if these had been already significant in Fig. C1. We saw no evidence of our strongest LyC candidates being located in the same physical CCD region of each exposure. On the contrary, our strongest LyC emitters are known “Galaxies with weak AGN” (Table 2), and show no correlation in their observed position on CCDs. On average, their LyC flux is ~ 1.0 mag brighter than that of our “Galaxies without AGN”. The AGN LyC flux is sometimes compact ($\lesssim$ a few pixels; Figs. 8 and B3) — as opposed to that of the galaxies (see Figs. 7 and B2) — and therefore occurs on too small of an optical scale to be caused by the severely out-of-focus redleak through pinholes. Thus, since the physical location of the AGN on the CCDs is spatially uncorrelated, their enhanced LyC flux is unlikely associated with residual sky-background gradients from pinholes. Had that been the case, a number of our much more numerous “Galaxies without AGN” would likely have also been exposed near pinhole-induced sky-gradients

and shown the same amount of red-leaked flux, which is not the case. Hence, local residual sky-background gradients due to pinhole-induced redleak enhancements are not likely to have affected our LyC measurements, at least not to $AB \simeq 32$ mag arcsec $^{-2}$.

B.3. Stacking Galaxies with Uncertain Redshifts

Cen & Kimm (2015) show with hierarchical simulations that the average LyC f_{esc} -values determined from stacks of galaxies will converge to more accurate and representative values with increasing sample size, simply because the average is better constrained from the increased sample statistics. When sample sizes are low, these average are more uncertain, and the flux measured from stacks also become more sensitive to contamination, since the contribution of flux from one galaxy is inversely proportional to the sample size. This sensitivity is augmented by the relative brightness between LyC and non-ionizing UVC in a stellar SED, where the UVC can be ~ 10 –40 $\times$ brighter than LyC in a stack, even slightly redward of the Lyman break. Thus, stacking flux from galaxies with some incorrectly determined redshifts can cause the total LyC signal to appear much brighter than what the galaxy SEDs would imply, as flux from the incorrect portions of the spectra of these galaxies is co-added to real LyC emission from others. However, stacking some contaminating flux with real LyC can be useful in verifying the stacking method and subsequent photometry in the stacks with better statistics, given that the real LyC signal is very faint, making its validation in the more reliable HQ stacks of fewer exposure more difficult. We therefore stacked the LyC and UVC emission of all galaxies with HQ and IQ (HQ+IQ) spectra, as well as separately for galaxies with and without AGN features, to ascertain how our stacking methods would be affected by larger sample statistics with the less reliable samples, and how this would affect the resulting LyC photometry.

For our larger, but likely more contaminated total sample of galaxies with HQ+IQ spectra, the corresponding WFC3/UVIS F225W stack contains co-added sub-image frames of 29 galaxies at $2.262 \leq z \leq 2.450$ with a mean redshift $\langle z \rangle = 2.362$, the F275W stack has 26 objects at $2.470 \leq z \leq 3.076$ with $\langle z \rangle = 2.669$, the F336W stack has 23 objects at $3.111 \leq z \leq 4.149$ with $\langle z \rangle = 3.505$, and the ACS/WFC F435W stack contains 37 objects at $4.379 \leq z \leq 6.277$ with $\langle z \rangle = 5.081$. The LyC and UVC stacks for these galaxies are shown in Fig. B1, in Fig. B2 for only the galaxies with *no* evidence for AGN activity, and in Fig. B3 for only galaxies with AGN signatures.

Table B1 shows that LyC flux levels corresponding to $m_{\text{AB}} \simeq 29.75, 29.53, 30.00,$ and 30.25 mag for F225W, F275W, F336W, and F435W at a measured formal SNR of $\sim 9.2, 11.6, 6.4,$ and 6.3 , respectively, when using the optimal apertures defined in the smoothed LyC stacks (blue ellipses). We find similar LyC fluxes ($m_{\text{AB}} \simeq 29.70, 29.37, 30.25,$ and 31.17 mag) with SNR $\sim 8.9, 11.8, 5.6,$ and 2.9 , respectively, when using the larger UVC apertures (green ellipses). The signal and corresponding noise values of the two measurements are generally consistent, indicating that our methods are in fact correctly averaging the central LyC flux and sur-




Figure BB1: Sub-image stacks of galaxies in the four different redshift bins with HQ+IQ spectra, sampling LyC emission in: [a] F225W at $2.26 \leq z \leq 2.45$, [b] F275W at $2.47 \leq z \leq 3.08$, [c] F336W at $3.11 \leq z \leq 4.15$, and [d] F435W at $4.38 \leq z \lesssim 6.28$; and corresponding UVC emission in: [e] F606W, [f] F606W, [g] F775W, and [h] F850LP.




Figure BB2: As Fig. B1 for only the galaxies without AGN with HQ and IQ spectra.




Figure BB3: As Fig. B1 for only the galaxies with AGN signatures and HQ and IQ spectra.

rounding backgrounds, and that our photometric measurements do indeed reflect the true value of the average LyC flux from the stack.

We note that the stacked LyC-fluxes of our sample of galaxies with HQ+IQ are significantly brighter than the stack of the galaxies with HQ spectra: $\Delta m_{\text{AB}}[\text{HQ+IQ}-\text{HQ}] \simeq 0.12 \pm 0.13$ mag (on average) when comparing the blue apertures and 0.23 ± 0.15 mag for the green apertures. This suggests that our HQ+IQ stack may not be significantly contaminated by non-ionizing flux from galaxies with incorrectly determined redshifts, since it gives similar LyC fluxes as the smaller and most reliable HQ stack, and so the fraction of incorrect redshifts of the galaxies with HQ+IQ spectra may still be low.

This also suggests that the size of our reliable $N \simeq 10\text{--}15$ objects in our HQ spectroscopic samples may just be large enough to yield average LyC-fluxes (and f_{esc} values) that are representative for the populations at those redshifts. The trend implied by Cen & Kimm (2015) is seen in our data: for our reliable $N \simeq 11\text{--}15$ galaxies, the measured SNR obtained on the average LyC-flux is less ($\text{SNR} \gtrsim 3\text{--}4$ for the blue apertures) than that obtained for $N \simeq 23\text{--}37$ HQ+IQ galaxies ($\text{SNR} \gtrsim 5\text{--}6$ for the same apertures). That is, the *average LyC-flux* for the larger samples used in our HQ+IQ stacks may indeed be more accurately determined than in our smaller HQ galaxy samples alone, even though both samples yield, to within their uncertainties, comparable average LyC fluxes. This implies that our photometric measurements are indeed determined correctly, and that the photometry of our stacks of galaxies with reliable redshifts may be near the true average value for galaxies at their respective redshifts, masses/luminosities, SFRs, A_V -values, ages, etc.

C. SOURCES OF SYSTEMATIC UNCERTAINTIES

C.1. Impact of Gradients in the Residual Sky-Background

Subtle gradients still exist in the new ERS UV mosaics, but at a much reduced level from the v0.7 ERS mosaics of W11. These are now $\lesssim 3\text{--}5\%$ of the Zodiacal sky-values from *corner to corner* across each of the eight individual 4096×4096 pixel CCD images that were drizzled onto the UVIS mosaics. After rebinning to $0''.09$ pixels, each of the individual WFC3 CCD exposures becomes ~ 1800 pixels in size when they are being co-added to the mosaics. This subtle gradient pattern was not very discernible, but appears to be similar in each of the 8 full WFC3/UVIS CCD frames in the ERS to a good approximation, and roughly linear across each CCD. The cause of these remaining gradients could be subtle residual errors in the on-orbit master bias-frames, in the delta-flat corrections used in the recent WFC3 pipeline reduction, and/or from variation in exposure time or background noise across the drizzled mosaic (Baggett & Anderson 2012; Mack et al. 2013). These remaining gradients are too faint to accurately map and remove from individual WFC3/UVIS UV exposures prior to drizzling, and removal of inaccurately measured gradients would introduce additional unintended errors in the mosaics. We therefore will assess the effects that these $3\text{--}5\%$ global gradients have on the $\lesssim 71$ pixel scales at which local sky-subtraction is performed in the LyC image stacks.

Dividing each LyC stack into our 9 segment “tic-tac-toe” grid, we determine the sky-background level and uncertainty in each of the 8 segments around the central box that contains the LyC candidate itself, which we exclude from the sky-background calculation. We compute these by optimally binning the count rates of the 8 outer segments, then we fit a normal distribution to the inner quartile of this data, taking the average of the fitted distribution as the sky-value and the $+1\sigma$ value to be the 84% of the pixel histogram. We estimate the gradients in the stacks from the rms value of the fitted average count rate in each segment. As shown in the examples of our “tic-tac-toe” photometric analysis in Fig. A3 and tabulated in Table A1, we find that any residual sky-background *gradient* left in the image stacks is $\sim 5.2\text{--}40 \times$ ($\sim 1.8\text{--}4$ mag) fainter than the residual sky-background numbers derived from Fig. 3. This then implies that *any* gradients in the local sky-background in the LyC image stacks (containing 10–37 objects each) are fainter than $\sim 32.3, 32.1, 32.5$, and 32.0 mag arcsec $^{-2}$ in WFC3/UVIS F225W, F275, and F336W, and ACS/WFC F435W across the $6''.39 \times 6''.39$ extent of each sub-stack, respectively.

These numbers are consistent with the aforementioned $\sim 3\text{--}5\%$ linear gradient across each of the full WFC3/UVIS mosaic images *before drizzling*, and corresponds to a $\lesssim 0.2\%$ error in the sky-subtraction across typical 71×71 pixel sub-image stacks. For a UV sky-brightness of $\mu_{\text{sky}} \sim 25.5$ mag arcsec $^{-2}$ (see Table A1), this amounts to a sky-subtraction error of ~ 32.3 mag arcsec $^{-2}$ across a 71×71 pixel stack. One possible source of such gradients are residual dark-current subtraction errors. Rafelski et al. (2015) show that the 2009 WFC3/UVIS dark current may vary between $0.00045 e^-/\text{s}$ and $0.00035 e^-/\text{s}$ across the CCDs (black curve in their Fig. 15). From experience, the quality of the calibration files is such that these gradients are typically subtracted at the level of (conservatively) $\sim 20\%$ of the gradient itself. That is, this dark current subtraction error across the 71 rebinned-pixel “tic-tac-toe” sub-images (out of 1800 rebinned pixels across the two CCDs) will amount to a residual sky subtraction error in the subimages of approximately: $-2.5 \log [0.20 \times ((0.00045/0.00035) - 1) * 71/1800] = 6.62$ mag below sky. This could then leave a residual dark current gradient on top of the UV zodiacal sky (25.5 mag arcsec $^{-2}$) of $25.5 + 6.6 = 32.1$ mag arcsec $^{-2}$, consistent with the limits given above. This level of uncertainty in the local sky-background level may pose a fundamental limit on the sensitivity and accuracy of any LyC (surface) photometry, which is slightly fainter than that potentially imposed by pinholes. The systematic nature of these residual gradients also explains the slight improvement in SNR noted in Appendix A.1 when the individual F225W sub-images are rotated by random multiples of 90° . These residual gradients are also much fainter than the measured LyC signal in general.

C.2. Assessment of Possible WFC3/UVIS CTE degradation $\lesssim 4$ Months After Launch

Charge Transfer Efficiency (CTE) degrades over time due to high energy cosmic ray collisions with the detector, and from encounters with relativistic protons and electrons during

Table BB1.: LyC Stack Photometry

Filter (1)	z-range (2)	$\langle z \rangle$ (3)	N_{obj} (4)	LyC apertures				UVC apertures				
				m_{LyC} (5)	$\text{ABerr}_{\text{LyC}}$ (6)	SNR_{LyC} (7)	A_{LyC} (8)	m_{LyC} (9)	SNR_{LyC} (10)	A_{UVC} (11)	m_{UVC} (12)	SNR_{UVC} (13)
ALL GALAXIES:												
F225W	2.262–2.450	2.362	33	29.93	0.12	9.30	0.935	29.75	9.19	1.364	24.62	314.7
F275W	2.470–3.076	2.669	33	29.60	0.08	13.94	0.797	29.27	14.46	1.386	24.81	268.8
F336W	3.111–4.149	3.505	27	29.99	0.15	7.23	0.988	29.96	7.34	1.008	24.69	193.8
F435W	4.379–6.277	5.263	40	29.84	0.15	7.35	0.660	30.24	5.62	0.559	26.27	109.7
GALAXIES WITHOUT AGN:												
F225W	2.262–2.450	2.363	29	29.75	0.12	9.21	1.137	29.70	8.69	1.445	24.54	307.6
F275W	2.481–3.076	2.662	26	29.53	0.09	11.61	1.095	29.37	11.84	1.395	24.78	242.0
F336W	3.111–4.149	3.536	23	30.00	0.17	6.42	1.031	30.25	5.57	0.881	24.76	174.7
F435W	4.414–6.277	5.312	37	30.25	0.17	6.27	0.448	31.17	2.93	0.396	26.68	90.3
GALAXIES WITH AGN:												
F225W	2.291–2.448	2.360	4	30.10	0.26	4.17	0.347	30.53	1.41	1.563	25.33	266.6
F275W	2.470–3.009	2.697	7	29.23	0.13	8.29	0.837	29.60	6.91	0.618	25.02	159.2
F336W	3.217–3.474	3.326	4	29.53	0.21	5.17	0.612	29.34	5.47	0.777	24.52	102.9
F435W	4.379–4.823	4.654	3	28.09	0.12	9.08	0.351	28.09	8.07	0.468	24.75	80.3

Table columns: As for Table 2, but for galaxies with HQ+IQ spectra (see Figs. B1–B3).

HST’s frequent passages through the South Atlantic Anomaly. Particle damage to the silicon of the CCD can cause areas where electrons become trapped in the detector’s crystal lattice during readout of the array. The WFC3/UVIS detectors suffer from a CTE loss of ~ 0.1 mag per year. After several years in orbit, faint objects ($\lesssim 300e^-$) can lose up to 50% of their flux during readout (Noeske et al. 2012; Bourque & Kozhurina-Platais 2013). CTE degradation can also cause charge trails to be visible in the images, caused by the delayed release of trapped electrons during readout. Partial recovery and correction of CTE losses in post-processing of the images is only possible for brighter sources (Anderson & Bedin 2010; Massey et al. 2014). Flux from very faint objects cannot be corrected in this manner, as their low electron counts are lost in the background noise of the detector. Because the WFC3 UV data were taken less than four months after Shuttle Servicing Mission SM4 that installed WFC3 onto *HST*, the WFC3/UVIS ERS data may not yet suffer from significant CTE losses. Similarly, the GOODS ACS/WFC F435W data, collected within five months of the SM3B repair mission in March 2002, may not be significantly affected by CTE degradation either. The data in the F435W filter also have much higher zodiacal sky-background, which provides sufficient count levels to fill any charge traps in the chips.

To be certain that the effects from CTE-degradation are still low enough for our LyC study with the ERS data, we also created two separate sub-stacks for each filter that were subdivided by their total expected *cumulative* CTE in the parallel-readout (or Y) direction of the UVIS CCDs. The CTE may be lower in some regions further away from the amplifiers at the beginning of readout due to any existing electron traps first being filled nearest to the amps during readout. However, this effect is very non-linear, as the traps have an unknown probability of capturing and releasing electrons per each charge

transfer, so electrons further away from the amps might encounter more traps if the captured electrons are released before the last columns are read out. Thus, the *cumulative* CTE will be higher on average closer to the amps when combining all CTE effects along the *parallel* readout direction.

Fig. C1a provides an overview of the WFC3/UVIS readout configuration (see Fig. 6.14 of Dressel et al. 2015). Fig. C1b shows the same for ACS/WFC F435W, which has identical readout directions to WFC3/UVIS, but its detectors have slightly different dimensions (2048×4096 pixels for ACS/WFC vs. 2051×4096 for WFC3/UVIS). The two $2k \times 4k$ WFC3/UVIS CCDs are shown as rectangular panels in Fig. C1, displaying the total pixel arrays from $X=1-4096$ and $Y=1-2051$ pixels. Within each WFC3/UVIS chip in Fig. C1, the green band indicates the half of each CCD *closest* in the parallel read-out direction to the four corresponding amplifiers (labeled as Amp A, B, C, D, respectively), where the cumulative CTE degradation effects from CR and relativistic particle hits accumulated on-orbit are expected to be *lowest*. We refer to this higher CTE region as the “high-CTE” area. The red bands in each chip indicate the half of each WFC3 CCD *furthest* from the amplifier in the Y read-out direction, where the effects from CTE-degradation would be more substantial. The cumulative CTE value itself from these areas will be lower compared to the “high-CTE” regions, since its signal will have on average been read-out through more charge transfer rows. We refer to this region as the “lower-CTE” area.

The colored circles in Fig. C1ab show, in *physical* WFC3/UVIS CCD (X,Y)-coordinates, where all our galaxies at $z \simeq 2.3-6$ were observed in all individual exposures within each of the two CCDs for both WFC3/UVIS and ACS/WFC, respectively. The filled circles indicate where some LyC flux was visible by eye for that individual galaxy in the respective




Figure CC1: Detector layout and observed location of galaxy sub-samples. (a) WFC3/UVIS F225W, F275W, and F336W images. The two $2k \times 4k$ CCDs are shown as the rectangular panels that stretch from $X=1$ to 4096 and $Y=1$ to 2051 pixels. Within each WFC3 CCD, the green regions indicate the half of each CCD quadrant that is closest to the corresponding read-out amplifier along the parallel read-out direction, while the red regions indicate the half that is furthest away and will undergo more transfers along the array. The colored circles mark the positions in physical WFC3/UVIS CCD coordinates of each of the $z \simeq 2.3\text{--}4.8$ galaxies with HQ spectra in our sample. Filled circles mark objects where visible flux is seen in the individual LyC sub-images. The sub-samples of objects in the green and red regions are referred to as our “high-CTE” and “lower-CTE” sub-samples, respectively. (b) as (a) for the ACS/WFC F435W images within the ERS footprint of the GOODS-S observations. Averaged over all objects, the difference between the stacked LyC signal in the “high-CTE” are compared to the “lower-CTE” area is $\Delta(\text{High-CTE-Lower-CTE}) \simeq +0.46 \pm 0.19$ mag. This suggests that the CTE-induced systematics are not yet larger than the rms error and other systematics in the photometry. Since the filled circles are fairly uniformly distributed across the CCD, CTE-degradation must have not yet been a major limitation in detecting the faint LyC signal four months after the WFC3 installation, and less than a year after the ACS launch.

Table CC1: Assessment of the Impact of CTE Effects for Galaxies with HQ+IQ spectra

Filter	z -range	$\langle z \rangle$	N_{obj}	LyC apertures			UVC apertures				
				m_{LyC}	SNR_{LyC}	A_{LyC}	m_{LyC}	SNR_{LyC}	A_{LyC}	m_{UVC}	SNR_{UVC}
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
ALL GALAXIES, HIGH-CTE AREA:											
F225W	2.262–2.450	2.364	24	30.53	6.35	0.59	30.10	5.63	1.64	24.69	247
F275W	2.470–3.076	2.652	19	30.69	7.12	0.30	29.75	8.59	1.12	24.82	241
F336W	3.117–4.149	3.515	17	29.75	6.65	1.32	30.13	5.78	0.85	24.61	187
F435W	4.597–6.277	5.200	19	30.14	4.73	0.51	30.81	3.12	0.33	26.16	103
ALL GALAXIES, LOWER-CTE AREA:											
F225W	2.262–2.450	2.357	24	29.75	8.91	1.17	29.68	9.09	1.26	24.63	305
F275W	2.481–3.008	2.672	19	29.18	15.2	1.03	29.11	14.7	1.28	24.81	219
F336W	3.110–3.917	3.441	15	30.06	6.42	0.68	30.17	5.00	0.88	24.82	149
F435W	4.379–6.095	5.294	22	29.66	7.60	0.51	29.99	6.04	0.42	26.56	64

Table columns are as for Table 2.

filter mosaic, and empty circles indicate no visible LyC emission. Only galaxies with HQ spectra were plotted, including those with AGN. We note that the galaxy sample was specifically selected to have high reliability in spectroscopic redshifts (see §3.1 and the HQ sample in Tables 2 and 3), therefore any apparent spatial correlation of the objects in the CCD coordinates is not due to CTE-degradation, but from the spectroscopic object selection.

There are four drizzled exposures in both WFC3/UVIS F225W and F275W, and three exposures in WFC3/UVIS F336W. The ACS/WFC F435W mosaics each have six exposures in total: two unshifted exposures per orbit, where each of the three orbits yielded one independent drizzle point. Every object with a spectroscopic redshift to $AB \lesssim 25$ mag is therefore plotted up to $11\times$ in the WFC3/UVIS in Fig. C1a and $3\times$ for ACS/WFC in Fig. C1b to monitor their actual locations in the individual WFC3/UVIS F225W, F275W, F336W, and ACS/WFC F435W exposures, respectively. The somewhat apparent clustering of ERS galaxies with the most reliable redshifts in Fig. C1 is caused by this repetitive plotting of the individual locations of the galaxies in each exposure, overemphasizing the appearance of any real clustering in the mosaic. It is possible that the way the VLT and other spectroscopic masks were configured to observe galaxies for redshift measurements — while maximally avoiding spectral overlap — further introduced some apparent clustering in a particular region of Fig. C1. Since the VLT spectroscopic masks are comparable in size to the WFC3 FOV, it is possible that the objects who received the most integration time during spectroscopy were more preferentially selected to be on one side of the field-of-view to minimize the spectral overlap.

With these two separate regions subdivided by their aggregate CTE, we can now determine if and how much the LyC measurements vary based on their location within each chip, and if CTE was already significantly degraded in the September 2009 WFC3/UVIS mosaics less than four months after WFC3's launch. Similar to our randomly subdivided test stacks in Appendix A.1, we combined these two sub-samples in each filter to compare their photometry, background, and noise levels. Some objects appear both in the “high-CTE” and “lower-CTE” areas of Fig. C1ab, which occurred due to the dithering of the individual WFC3/UVIS ERS exposures. These objects were added to the corresponding sub-stacks for assessing the effects from CTE-degradation. Fig. C1ab show that about as many objects with spectroscopic redshifts are located in the green (“high-CTE”) areas as in the red (“lower-CTE”) areas, allowing us to make a quantitative comparison between the two subsets to assess the possible effects of differential CTE across the detector.

The results for the two HQ+IQ CTE sub-samples are shown in the bottom two sets of rows in Table C1. From these measurements, as a function of relative position on the CCDs, it is clear that each of these two CTE sub-samples still yields significant LyC detections, with correspondingly larger errors due to the smaller number of objects in each sub-sample. In particular, the LyC flux in the “lower-CTE” sub-sample remains at least as significant as it is for the “high-CTE” sub-sample: on average, the LyC flux in the “lower-CTE” sub-

sample is 0.46 ± 0.19 mag (rms) brighter than in the “high-CTE” sub-sample (Table 2). This difference in photometry between the “high-CTE” and “lower-CTE” sub-samples — as averaged over the four filters — is marginally significant, but its sign is such that the “lower-CTE” sub-sample is actually somewhat brighter than the “high-CTE” sub-sample. *Therefore this difference does not point to significant WFC3 CTE effects as of Sept. 2009.* The difference also varies fairly randomly between the four filters, showing no general trend with Y-position on the CCD's. Small differences in the observed LyC flux from the two different regions on each CCD may also be caused by some intrinsic variation of LyC f_{esc} between the two sub-samples, and/or by the small number statistics available in both sub-samples in general. Thus, any CTE induced systematics four months after WFC3's launch are not yet larger than the random errors in the faint LyC signal. This is consistent with the decline in CTE of ~ 0.1 mag/year measured by (Noeske et al. 2012; Bourque & Kozhurina-Platais 2013).

In Fig. C1a, filled circles indicate where marginal LyC signal is marginally visible in the individual sub-images. This LyC signal is generally no brighter than $2-3\sigma$ spread over a few pixels for an individual object, so we do not further analyze the LyC signal from individual objects here. This information, however, does allow us to do a simple statistical analysis to see if CTE-degradation has already affected the fraction of objects with marginally visible LyC flux between the “high-CTE” and “lower-CTE” areas of the WFC3 CCDs. For this, we used the total of 14 AGN and 50 galaxies with HQ spectra and no AGN (Table 2) covered by WFC3 UVIS and ACS/WFC images in the filters F225W, F275W, F336W, and F435W, respectively. Since these were taken in 4, 4, 3, and 3 dither points each, for WFC3 this resulted in $4 \times 1 + 4 \times 7 + 3 \times 4 = 44$ circles in Fig. C1a for AGN, and $4 \times 14 + 4 \times 11 + 3 \times 10 = 130$ circles for HQ galaxies, or a total of 174 circles. Drizzling moved 3 of these off the CCD-area, so in total 171 circles are visible in Fig. C1a, 100 of which are in the “high-CTE” area, and 71 in the “lower-CTE” area.

Of the total of 171 objects observed on a UVIS CCD at any time in any filter in Fig. C1a, a total of 65 or $\sim 38\%$ show marginal LyC signal in the individual images. Of these marginal detections, 35/100 or $\sim 35\%$ occurred in the “high-CTE” area and 30/71 or 42% occurred in the “lower-CTE” area. Since the filled circles are fairly uniformly distributed across the both individual WFC3 CCDs, and visible LyC detections exist in regions furthest away from the readout amplifiers, CTE must have been high enough for these faint detections to survive the multiple transfers during readout without becoming trapped in the detector. A similar conclusion is implicit from Fig. C1b for the ACS data, but with fewer statistics for the galaxies with HQ spectra covered in ACS F435W. Hence, neither the WFC3 UVIS or ACS data suffered from CTE effects that affected the possible fraction of LyC detections within 4–5 months after launch for WFC3 and ACS, respectively.

Both of these tests suggest that, less than four months after WFC3's launch, CTE-degradation in the UVIS chip is not

yet at a level that significantly affects the readout and subsequent photometry of faint flux to within the errors of our measurements, nor has it biased the distribution of objects with marginal individual LyC detections across the CCD. If this had been the case, we would have seen significantly larger differences in LyC flux across the two CCD detectors in the parallel readout direction, and would in fact have seen faint objects fully disappear into the noise in the “lower-CTE” regions in Fig. C1, in addition to visible charge trails for all brighter objects, none of which appear in our data.

D. MODELING AND UNCERTAINTIES

D.1. SED Fitting Uncertainties due to Extinction

In order to obtain accurate estimates for LyC and UVC dust extinction and subsequent absolute LyC escape fractions, we must adopt the best available dust attenuation models for galaxies with accurate redshifts and no contaminating AGN signatures when performing the minimized χ^2 stellar SED fits that results in the most likely A_V -values from the observed panchromatic ERS data. From SED fitting of Calzetti et al. (2000) attenuated BC03 models using the 10-band WFC3+ACS photometry of all ~ 6900 galaxies at $2 \lesssim z \lesssim 6$ within the GOODS-S ERS field (Windhorst & Cohen 2010; W11, see also §5.1), we find that most ERS galaxies have $0.0 \lesssim A_V \lesssim 1.0$ mag, although a few have best-fit extinctions as high as $A_V \simeq 2\text{--}2.5$ mag.

Fig. D1a shows the distribution of dust extinction A_V -values from best-fit SEDs for all galaxies in the 10-band ERS data (small black dots), compared to our spectroscopic sample in the four indicated redshift bins. The SED fitting sampled the A_V parameter space in 0.2 mag intervals. To facilitate plotting, a random fraction of 0.1 mag was added in quadrature to each best fit A_V -value in Fig. D1a. Fig. D1b shows the resulting $N(A_V)$ distributions from the SED fits of the galaxies with HQ+IQ spectra (top panel), as well as in each of our four redshift bins (bottom four panels). The best fit median A_V -values are indicated for each, as well as their $\pm 1\sigma$ ranges. These are listed for Galaxies with and without AGN. Because the $N(A_V)$ distribution contains a significant population of objects with A_V close to ~ 0 mag at all redshifts, and possibly an increasing fraction of more dusty objects towards lower redshifts, *no single exponential* in A_V/A_V^{med} describes the full $N(A_V)$ distribution in Fig. D1a and D1b over all redshifts. Nonetheless, because an analytical form of the $N(A_V)$ as a function of redshift is useful for some of our calculations, we used the generalized exponential of Windhorst et al. (1990) to describe $N(A_V, z)$ more fully as following:

$$N(A_V, z) = N_0 \cdot \exp(-\ln(2) \cdot (A_V/A_{V,\text{med}(z)})^n) \quad (\text{D1})$$

where N_0 is the number of objects in each redshift bin, and the exponent n is close to unity, but slowly varying with redshift to best describe the observed $N(A_V, z)$ distribution. Table 3 lists the A_V^{med} values and their $\pm 1\sigma$ ranges a function of redshift.

Fig. D1a shows that the A_V -values of our galaxy samples with spectroscopic redshifts are consistent with those found for the entire ERS sample of 6900 galaxies with 10-band fit-

ted photometric redshifts to $m_{\text{AB}} \lesssim 27$ mag. This implies that our galaxies *are* sampling the available parameter space of A_V -values at their approximate redshifts. We can thus quantify representative values of A_V^{med} , as well as its $\pm 1\sigma$ ranges for our galaxy populations as a function of redshift. Our calculation of the errors of the “direct” f_{esc} values (see §5.1) can then include this rms error, since the A_V -uncertainty in the SED fits is unknown, but would not be larger than the sample variance. The MC simulated $f_{\text{esc}}^{\text{abs}}$ values in §5.1 and §5.2 do not include an A_V uncertainty, even though they utilize the extinction-corrected (intrinsic) SEDs. This implicit A_V -error is one of the dominant errors in the “direct” f_{esc} calculation, but is less important than the IGM transmission variations in the MC derived f_{esc} values (see §5.2 and §5.3). We also note that there may exist a degeneracy between the A_V -values and the ages of the best fit SEDs, which would also add to the uncertainty of the A_V -values. The A_V -induced error can be as large as the combined uncertainty of the photometric observations that we fit, and can further increase the overall uncertainty of the SED fit. Furthermore, the error in R_V derived by Calzetti et al. (2000) ($R_V = 4.05 \pm 0.80$) is not propagated into the SED fit, which would also increase the A_V -error. However, the uncertainty in IGM transmission, which is primarily due to variations in sight-lines and redshifts in the stacks (see Inoue & Iwata 2008), dominates the error in the MC f_{esc} values, and so an additional A_V -error would only slightly increase their $\pm 1\sigma$ values, as discussed in §5.2 and Table 3.

Table 3 also shows that A_V^{med} , for the samples with *reliable* spectroscopic redshifts, increases from $\sim 0.17\text{--}0.18$ mag at $z \sim 3\text{--}5$ to $\sim 0.55\text{--}0.58$ mag at $z \sim 2.3\text{--}3$, consistent with the behavior seen in the entire ERS sample as a gradual increase in reddening towards the lower redshifts, when the stellar populations have aged more and produced more dust over cosmic time. Hence, the median A_V -values and their rms for galaxies appears to increase at the lower redshifts, and so our “direct” f_{esc} -error calculations also incorporate this sample variance as well.

We then converted A_V to A_{FUV} in order to be able to correct for the effects from dust in the UVC, so that the $F_{\nu,\text{UVC}}^{\text{int}}$ and $F_{\nu,\text{UVC}}^{\text{obs}}$ terms cancel in the “direct” calculation of f_{esc} (Eq. 5) to give us the escape fraction of LyC only. For star-forming galaxies, we use: $A_{FUV} \simeq 7.9 E_{B-V}$ mag, following Gil de Paz et al. (2007) or $A_{FUV} \simeq 10.33 E_{B-V}$ mag (Calzetti et al. 2000). Since $A_V \simeq 3.1 E_{B-V}$, we will adopt here: $A_{FUV} \simeq 2.55 A_V$ mag for Gil de Paz et al. (2007) dust, or $A_{FUV} \simeq 3.33 A_V$ mag for Calzetti et al. (2000) dust, respectively. We then can use these A_{FUV} values to convert our “direct” f_{esc} values as calculated in §5.1 to the dust-corrected f_{esc} , or a more uncertain $f_{\text{esc}}^{\text{abs}}$ value, as indicated in 3b.

D.2. LyC and UVC Surface Brightness Model Details

For the UVC and LyC SB-models in §5.5 and Fig. 11, we generate the UVC surface brightness as following:

$$SB_{\text{UVC}}(b) = \int_{-\infty}^{\infty} ds \epsilon_{\text{UVC}}(b, s). \quad (\text{D2})$$




Figure DD1: [Left]: Distribution of dust extinction A_V -values from best-fit Calzetti et al. (2000) dust attenuated BC03 SEDs for all galaxies in the 10-band ERS data (black dots), compared to the spectroscopic samples used for the stacks in the four indicated redshift bins (colored open circles for galaxies and open asterisks galaxies harboring AGN). [Right]: Adopted distributions $N(A_V)$ for our population of galaxies with HQ and IQ spectroscopic redshifts (top panel), as well as in each of our four redshift bins (bottom four panels). The best fit median A_V -values are indicated for each, as well as their $\pm 1\sigma$ ranges adopted in the error calculation of our “direct” f_{esc} values.

where b denotes the impact parameter, s denotes the line-of-sight coordinate, and $\epsilon_{\text{UVC}}(b, s)$ denotes the emission rate of UVC photons per unit volume at (b, s) . The distance from the galaxy r is defined as $r = \sqrt{b^2 + s^2}$ (Note that $2rdr = 2sds$ and $ds = \frac{r}{\sqrt{r^2 - b^2}} dr$, since $r^2 = s^2 + b^2$). We then obtain the LyC surface brightness from:

$$SB_{\text{LyC}}(b) = \int_{-\infty}^{\infty} ds \epsilon_{\text{UVC}}(b, s) \frac{f_{\text{LyC}}}{f_{\text{UVC}}} f_{\text{esc}}(b, s). \quad (\text{D3})$$

where $f_{\text{esc}}(b, s)$ denotes the fraction of LyC photons that can escape from (b, s) , and the factor $\frac{f_{\text{LyC}}}{f_{\text{UVC}}}$ simply rescales the flux at UVC frequencies to that at LyC.

In our model of a clumpy ISM $f_c(r)$ denotes the number of self-shielding clumps per unit length at r (see Dijkstra & Kramer 2012). We therefore find that $f_c(r)dr$ denotes the number of self-shielding clumps along a differential length dr . We assume that each clump is optically thick to ionizing photons. In this case, the escape fraction from (b, s) is simply the probability of finding at least one clump on a sight-line to (b, s) , $P_{\text{clump}}(b, s)$, which is given by:

$$P_{\text{clump}}(b, s) = 1 - P_{\text{noclump}}(b, s) = 1 - \exp[-N_{\text{clump}}(b, s)], \quad (\text{D4})$$

where in the last step we assumed that the number of clumps along a given line-of-sight follows a Poisson distribution with mean $N_{\text{clump}}(b, s)$. This mean is given by:

$$N_{\text{clump}}(b, s) = \int_{-\infty}^s ds' f_c(b, s'). \quad (\text{D5})$$

Table DD1.: List of Individual Galaxies with HQ and IQ spectra

HQ SAMPLE							IQ SAMPLE						
RA (J2000.0) Dec	z	$m_{J,AB}$	$M_{1500\text{\AA}}^{AB}$	$(V-I)$	AGN?		RA (J2000.0) Dec	z	$m_{J,AB}$	$M_{1500\text{\AA}}^{AB}$	$(V-I)$	AGN?	
[deg]	[deg]	[mag]	[mag]	[mag]			[deg]	[deg]	[mag]	[mag]	[mag]		
53.0065856	-27.7241780	2.726	22.84	-20.98	0.16	yes	53.0049707	-27.7344196	2.561	23.67	-20.84	0.14	no
53.0076531	-27.7018322	3.603	24.99	...	0.01	no	53.0054590	-27.7122028	2.300	23.84	-20.13	0.13	no
53.0088555	-27.7243680	2.726	24.00	-19.94	0.64	yes	53.0062813	-27.7396132	2.512	24.84	-19.59	0.19	no
53.0126417	-27.7472499	2.573	23.78	-19.11	0.69	yes	53.0094224	-27.7362606	2.555	24.13	-20.44	0.14	no
53.0135070	-27.7552439	3.217	24.51	-21.23	-0.03	yes	53.0138291	-27.7573623	3.472	24.05	-21.36	0.27	no
53.0139300	-27.7568368	2.317	23.67	-21.35	-0.08	no	53.0162071	-27.7212277	2.852	24.40	-20.73	0.29	no
53.0145364	-27.7279355	3.132	24.26	-21.12	0.29	no	53.0202764	-27.7476192	3.404	23.47	-21.97	0.44	no
53.0205689	-27.7421565	3.474	23.55	-22.10	0.18	yes	53.0271513	-27.7164362	3.605	24.77	-21.02	0.49	no
53.0209195	-27.7701853	3.917	24.58	-21.57	0.63	no	53.0352291	-27.7441260	4.149	25.58	-20.72	0.56	no
53.0210399	-27.7704500	3.912	...	...	...	no	53.0378822	-27.7280828	2.299	25.64	-19.34	-0.14	no
53.0211680	-27.7823660	4.823	23.51	...	2.25	yes	53.0381127	-27.7658301	2.325	23.60	-20.60	0.17	no
53.0213294	-27.7504368	2.559	23.46	-20.93	0.17	no	53.0393892	-27.7662312	2.330	23.92	-20.51	0.09	no
53.0219086	-27.7167982	4.812	25.23	-21.24	2.03	no	53.0488132	-27.6971239	4.912	25.23	-21.30	1.70	no
53.0344441	-27.6982203	2.470	24.66	-19.65	0.08	yes	53.0491167	-27.6913003	3.338	23.62	-21.29	0.29	no
53.0357043	-27.7306428	2.365	24.53	-19.89	0.09	no	53.0528107	-27.6949474	2.718	25.22	-20.10	-0.01	no
53.0417672	-27.6923910	4.811	24.80	-21.60	1.77	no	53.0538906	-27.7476242	3.111	24.08	-20.93	0.26	no
53.0424530	-27.7378682	2.304	23.04	-21.16	0.10	no	53.0589207	-27.6978274	2.288	23.56	-20.63	0.23	no
53.0432703	-27.7651153	2.567	24.14	-20.95	0.10	no	53.0626005	-27.7025228	2.783	23.49	-21.66	0.19	no
53.0497074	-27.6992041	5.578	26.21	-19.60	2.27	no	53.0632582	-27.6996461	2.402	23.65	-19.23	0.52	yes
53.0598383	-27.7092196	3.009	25.66	-20.08	-0.09	yes	53.0662303	-27.6899867	5.574	25.10	-21.52	1.51	no
53.0624487	-27.7355965	2.673	24.05	-20.79	0.18	no	53.0669686	-27.7405325	4.136	24.50	-20.39	0.23	no
53.0650211	-27.7326889	2.649	24.39	-20.76	0.02	yes	53.0687298	-27.7315272	2.437	24.75	-19.99	-0.01	no
53.0652442	-27.7429043	2.616	23.53	-21.39	0.14	no	53.0741807	-27.9115382	6.277	...	...	1.83	no
53.0654173	-27.7542521	2.673	23.55	-19.82	0.29	no	53.0753649	-27.6869757	5.979	27.04	...	4.68	no
53.0708237	-27.6871523	4.414	24.74	...	1.34	no	53.0787941	-27.6937455	2.306	24.45	-20.08	-0.01	no
53.0740190	-27.7327531	2.619	25.69	...	-0.04	no	53.0792894	-27.6913746	2.435	23.12	-21.13	0.12	no
53.0752483	-27.7551833	3.706	24.36	-21.35	0.54	no	53.0860664	-27.7441787	2.691	24.91	-20.69	-0.03	no
53.0780196	-27.7310295	2.416	23.43	-20.82	0.16	no	53.0872436	-27.7295359	3.596	24.60	-21.16	0.37	no
53.0788083	-27.8840868	5.563	24.52	-22.14	2.45	no	53.0953319	-27.7909781	5.128	24.59	-21.77	1.64	no
53.0793720	-27.7416250	3.356	24.45	...	0.60	yes	53.0993282	-27.9198850	6.095	...	...	0.00	no
53.0887662	-27.6808873	5.292	25.29	-21.33	1.99	no	53.1008187	-27.7159897	2.298	23.18	-20.60	0.16	yes
53.0914387	-27.7274352	2.576	24.50	-20.67	0.13	no	53.1033062	-27.7996620	5.996	27.18	-19.82	1.43	no
53.0953630	-27.7558710	4.440	...	...	...	no	53.1037314	-27.7140967	2.262	23.97	-19.73	0.17	no
53.0953874	-27.6875256	3.357	25.01	...	0.22	no	53.1054735	-27.7585599	4.992	26.82	-19.61	1.61	no
53.1066753	-27.9301691	5.786	24.63	-22.05	1.72	no	53.1076063	-27.7139852	3.770	24.90	-20.89	0.72	no
53.1070229	-27.7182311	2.291	23.28	-18.77	0.52	yes	53.1123707	-27.7450151	2.560	24.02	-20.44	0.33	no
53.1105738	-27.7483624	2.727	24.25	...	0.08	yes	53.1174463	-27.8051784	5.940	26.15	-20.53	-1.23	no
53.1124455	-27.6926271	2.440	23.25	-21.44	0.06	no	53.1190141	-27.6821532	4.597	25.33	-20.90	1.84	no
53.1129923	-27.7455481	2.323	24.35	-19.68	0.06	no	53.1209335	-27.7094289	3.585	24.97	-20.96	0.30	no
53.1133300	-27.6977122	3.168	24.91	-20.62	0.11	no	53.1218483	-27.7163861	2.577	24.55	-19.40	0.14	no
53.1178213	-27.7343079	3.256	22.38	...	0.21	yes	53.1234359	-27.6827260	3.117	23.96	-21.69	0.34	no
53.1202648	-27.6923720	4.800	23.74	-21.35	1.42	no	53.1254020	-27.8493553	5.087	23.77	-24.22	0.67	no
53.1205187	-27.6911605	4.882	...	...	...	no	53.1274909	-27.7109506	2.689	...	...	...	no
53.1206044	-27.7365858	3.368	24.72	-21.17	0.21	no	53.1303090	-27.6959093	2.438	24.32	-20.36	0.07	no
53.1216385	-27.6729132	3.308	25.10	-20.75	0.09	no	53.1336927	-27.6934559	3.697	24.76	-21.08	0.64	no
53.1220340	-27.9387290	4.760	24.45	-21.36	1.63	yes	53.1345577	-27.6906495	2.330	23.08	-21.68	0.08	no
53.1317173	-27.6690080	3.076	23.80	-21.75	0.16	no	53.1375638	-27.7001019	2.448	23.41	-19.67	0.46	yes
53.1348330	-27.7133537	2.430	23.19	-21.16	0.18	no	53.1382958	-27.6636273	5.830	...	...	3.73	no
53.1387530	-27.7004642	2.450	23.37	-19.95	0.29	no	53.1445021	-27.7280735	2.276	24.58	-19.53	0.14	no
53.1454255	-27.6980031	2.313	23.84	-20.81	-0.01	no	53.1456119	-27.6852449	2.771	24.18	-21.23	0.09	no
53.1536740	-27.7177307	2.449	23.97	-19.59	0.35	no	53.1568625	-27.6627048	2.399	24.10	-20.67	0.35	no
53.1553015	-27.7192736	2.400	25.14	-20.23	-0.02	no	53.1622952	-27.7121259	4.379	23.11	-21.26	0.45	yes
53.1567717	-27.8395448	5.518	25.37	-21.17	3.14	no	53.1627506	-27.7607401	5.920	27.73	-19.19	...	no
53.1574353	-27.7090018	2.975	23.67	-21.57	0.21	no	53.1641631	-27.7098940	2.448	23.67	-20.02	0.17	no
53.1619654	-27.7226503	2.449	23.64	-21.08	0.14	no	53.1658924	-27.8827931	5.543	26.12	-20.50	2.41	no
53.1679957	-27.7113456	2.584	23.82	-21.20	0.08	no	53.1667066	-27.8041552	5.828	25.12	-21.57	2.37	no
53.1695277	-27.6578205	2.302	23.54	...	...	no	53.1671490	-27.7598378	4.773	25.37	-21.02	1.72	no
53.1789565	-27.6958494	3.466	24.88	-20.99	0.28	no	53.1700564	-27.6839724	2.580	24.57	-20.56	0.26	no
53.1828039	-27.7052559	2.368	24.40	-20.88	-0.05	no	53.1720409	-27.6840723	2.323	23.57	-20.46	0.18	no
53.1836370	-27.7076241	4.923	25.05	-21.51	2.09	no	53.1747855	-27.6948481	2.577	24.42	-20.33	0.14	no
53.1892618	-27.9106873	4.923	24.90	-21.50	2.33	no	53.1760436	-27.6658902	2.801	25.08	-19.39	...	no
53.1896098	-27.6952668	2.584	24.56	-19.74	0.22	no	53.1777269	-27.8274860	4.840	25.03	-21.37	1.78	no
53.1785568	-27.7017316	3.412	24.14	-21.48	0.39	no	53.1882585	-27.7193896	2.481	23.80	-20.55	0.34	no
53.1813846	-27.8219979	4.763	25.83	-20.62	1.87	no	53.1884479	-27.8193893	5.583	25.78	-20.88	3.27	no
53.1918227	-27.8249116	5.787	26.55	-20.22	1.90	no	53.2294990	-27.9040151	4.718	...	...	2.55	no
53.1934334	-27.9234492	6.082	...	...	0.28	no	53.2254025	-27.8210893	5.793	24.87	-21.66	4.80	no
							53.2381863	-27.8624909	4.765	24.64	-21.71	1.42	no